

Self-Constructing Continua

Doug McLellan — Spring 2021 Draft

Abstract

Recent interest in “potentialism,” which conceives mathematics as growing over time, suggests that we try formalizing the idea of a mathematical universe that constructs itself in a gradual way, with no work from outside nor any inexplicable novelty involved in the course of its growth. Taking *continuum* to mean the set of all reals in a standard inner model of set theory, we axiomatize the assertion that a family \mathcal{F} of distinct continua constructs itself in such a way; we then present a forcing construction that outputs candidates $\mathcal{F}(B, \zeta, G)$ to satisfy our axioms, where B is a boolean algebra, G is a generic ultrafilter on B , and ζ is a function that (with G) determines which of B 's subalgebras contribute continua to the family. We derive a key necessary condition that B must meet for $\mathcal{F}(B, \zeta, G)$ to satisfy the axioms. We show that neither the Cohen-real nor the random-real algebra meets this condition, mainly because their proper subalgebras C are *non-rigidly included*, meaning the algebra has nontrivial automorphisms that leave C fixed. Rigid boolean algebras avoid this problem, but the example we give fails to yield a model of the axioms for other reasons. We briefly discuss the most promising candidate B , a “noncommutative analog” of the random-real forcing algebra which is not rigid but does, we conjecture, have rigidly-included subalgebras.

1 Self-Construction Axioms

By the *constructive closure* $\mathbb{R}(X)$ of a set X of real numbers we mean all the real numbers that must exist given that X does, and given that the ZF axioms hold. Formally we define $\mathbb{R}(X)$ for any set X as the set of all reals in $L(X)$, the constructible hierarchy seeded with the transitive closure of $\{X\}$ and then built up with definable operations using Gödel's well-known procedure. For background on $L(X)$ — which underlies everything that follows — see the appendix, Section 11.

By a *continuum* we mean a set X of real numbers (henceforth construed as subsets of ω) that is constructively closed, meaning $\mathbb{R}(X) = X$. Note that $L(X)$ is the smallest standard inner ZF model having X as a member (see Lemma 3.2) and it follows that we could define a continuum equivalently as the set of all real numbers in some standard inner ZF model.

$L(\emptyset)$, usually just denoted L , is the constructible universe of sets. Its continuum $\mathbb{R}(\emptyset)$ is the least continuum. Gödel famously proved it consistent with ZF that this is the *only* continuum.

It is also consistent with ZF that we can arrange distinct continua so that each is the constructive closure of the smaller ones' union; this holds, for example, of the set of all continua in a Cohen forcing extension of L . Such an arrangement suggests the real numbers being constructed out of each other — growing over time, as it were — or, if the continua are not *linearly* ordered by inclusion, then growing over space-time, as *it* were.

The goal of this paper is to determine whether this growth metaphor can be pushed further, whether a set of continua might grow “organically” without the need for anyone to do any preliminary “arranging.” We formalize this idea in two axioms, the first of which is stated in terms of “self-collection.”

Definition. A set \mathcal{N} of continua *self-collects into* X if the following hold:

- (i) $X \notin \mathcal{N}$; (X is a new continuum)
- (ii) $X, Y \in \mathcal{N} \Rightarrow (\exists Z \in \mathcal{N})(X, Y \subseteq Z)$; (\mathcal{N} is directed)
- (iii) $\mathcal{N} \in L(X)$; (nothing beyond X is used to construct X)
- (iv) $(\neg \exists x \in X)(\bigcup \mathcal{N} \subseteq \mathbb{R}(x) \subset X)$; (\mathcal{N} is unbounded below X)
- (v) $X = \mathbb{R}(\bigcup \mathcal{N})$. (X is the constructive closure of \mathcal{N} 's union)

Axioms for a Self-Constructing Family \mathcal{F} of Continua:

Self-Collection: $(\forall X \in L(\mathcal{F}))(X \in \mathcal{F} \iff$
some $\mathcal{N} \subseteq \mathcal{F}$ self-collects into X , and $X = \mathbb{R}(x)$ for some real x).

Foundation: $(\forall X \in \mathcal{F})(\forall x \in X)(\exists Y \in \mathcal{F})$
($x \in Y \subseteq X$, and $(\forall Z \in \mathcal{F})(Z \subset Y \Rightarrow x \notin Z)$).

Intuitively, the Self-Collection axiom captures an idea of continua naturally forming a collection that requires vanishingly little work to constructively close; and Foundation keeps real numbers from slipping in inexplicably or unaccountably during \mathcal{F} 's growth.

It is easy to check that $\mathcal{F} = \{\mathbb{R}(\emptyset)\}$ is a self-constructing family. This paper's concern is whether any less trivial \mathcal{F} can satisfy our two axioms.

2 The $\mathcal{F}(B, \zeta, G)$ recipe for singly-generated continua

The technique of forcing is uniquely suited to produce candidates to satisfy the self-construction axioms, and we restrict our attention accordingly:

Stipulation 2.1 *Henceforth every family \mathcal{F} we consider as a candidate to satisfy the self-construction axioms will belong to some forcing extension $L[G]$.*

We remark that for sets $Z \subseteq L$ (such as G), $L(Z) = L[Z]$.¹

The clause “ $X = \mathbb{R}(x)$ for some real x ” in the Self-Collection axiom amounts to another stipulation on \mathcal{F} , that every continuum in a self-constructing family be “singly generated”:

Definition. A *singly-generated* continuum X is one that satisfies $X = \mathbb{R}(x)$ for some real number x (which is clearly never unique).

The main reason for including the singly-generated clause in our axiom is that, when taken together with Stipulation 2.1, it gives our continua a particularly simple form. Namely, each will have form $\mathbb{R}(G \cap C)$, where C is a countably-completely-generated subalgebra of the boolean algebra on which G is a filter. (This is proved in Lemma 3.8; it is closely related to Lemma 15.43 of [6], the well-known “intermediate model theorem.”) Consequently a fully general recipe for a family of singly-generated continua in a forcing extension of L is given by

$$\mathcal{F}(B, \zeta, G) \equiv \{\mathbb{R}(G \cap C) : C \in \text{CSAs}(B) \text{ and } \zeta(C) \in G\},$$

where B is a complete boolean algebra in L , $\text{CSAs}(B)$ is the set of B 's countably-completely-generated subalgebras, G is a generic ultrafilter on B , and $\zeta : \text{CSAs}(B) \rightarrow B$ is a mapping in L that (along with G) picks out which subalgebras will contribute continua to the family.

The task of obtaining a self-constructing family thus reduces to finding a boolean algebra B and mapping ζ such that $\mathcal{F}(B, \zeta, G)$ satisfies our two axioms.

Our Foundation axiom, which requires for each real $x \in \bigcup \mathcal{F}$ that there be a continuum that is *least* in \mathcal{F} for having x as a member (or else that there be “enough continua *minimal* in \mathcal{F} for having x ,” in a certain sense) is not hard to satisfy. For example, if ζ maps every subalgebra to $1 \in B$, then each real $x \in L[G]$ does appear at a least member of $\mathcal{F}(B, \zeta, G)$, namely $\mathbb{R}(x)$.

The Self-Collection axiom, on the other hand, is apparently quite hard to satisfy. Our main question in this paper will be: When $\dot{\mathcal{N}}$ is a B -name for a subset of $\mathcal{F}(B, \zeta, G)$, under what conditions on B and $\dot{\mathcal{N}}$ will the latter be forced to self-collect into $\mathbb{R}(G)$?

¹ $\mathbb{R}(X)$ is defined with the “parentheses” version of relative constructibility ([6], Definition 13.24). A forcing extension $L[G]$ is defined differently, as the class of interpretations that the generic filter G gives to specified forcing names in L . This generalizes to a “square-brackets” version of relative constructibility ([6], Definition 13.22) which guarantees for arbitrary sets Z that $L[Z]$ satisfy the Axiom of Choice, but not that Z actually be a member of $L[Z]$. In particular, if \mathcal{N} is any set of continua then $L[\bigcup \mathcal{N}] = L$, hence $\mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(\emptyset)$, so the square-brackets version could not be used in clause (v) of self-collection. The square-brackets version is, however, the standard one to use when working with forcing extensions, so it is worth pointing out that when $Z \subseteq L$, e.g. when Z is a single real number or a filter on a constructible poset, $L(Z) = L[Z]$.

3 Main Definitions, Facts, Questions

We now turn to the definitions needed to answer the question just stated, and also to pose tighter versions of it that incorporate some simplifying assumptions. We will register these versions as our Main Question at the end of this section.

3.1 Definitions and Notation

$L(y)$ is the union of all levels $L_\alpha(y)$ of the (relative) constructive hierarchy seeded with $\{y\}$'s transitive closure (that is, $L_0(y) \equiv \text{TrCl}(\{y\})$); see Section 11 for details and background.

When $x \in L(y)$ we say that y *constructs* x . If also $y \in L(x)$, we say x and y are *interconstructible*.

We will not restate the basic terminology of forcing. We assume familiarity with the boolean-algebraic approach to forcing used in T. Jech's standard text [6], as well as in J. Bell's text [2]. We side with [2] in preferring the symbols \wedge and \vee for boolean meet and join, whereas [6] generally uses \cdot and $+$.

In what follows, B is always an atomless complete boolean algebra in L , the constructible universe. (We should say that the structure $(B, 0, 1, \vee, \wedge, \neg)$ is our boolean algebra and that B is just its underlying set, but we will conflate them as is customary.) We also assume B is countably completely generated (see the definition just below).

An *ACSA* of B is an atomless complete subalgebra of B . Switching “atomless” and “complete” would yield the more attractive acronym “CASA”, but “complete subalgebra of B ” is a syntactic unit: it denotes not just any subalgebra C of B that is complete, but only those C wherein the supremum of any subset of C is the same whether calculated in C or in B , and likewise for infima. See [6], chapter 7, under “Complete and Regular Subalgebras.”

If $X \subseteq B$, X *completely generates* the subalgebra \overline{X} that is the smallest complete subalgebra of B that includes X .

$\text{ACSAs}(B)$ denotes $\{C \in L : C \text{ is a countably-completely-generated ACSA of } B\}$, where by *countably-completely-generated* we mean that $C = \overline{Q}$ for some countable $Q \subseteq C$. As mentioned, we assume that B itself is countably-completely-generated.

C, D, E will always range over $\text{ACSAs}(B)$.

B^+ means $B \setminus \{0\}$, that is, B without its least element.

G is an L -generic ultrafilter on B (equivalently, L -generic filter on B^+).

$C \leq_G D$ means $(G \cap D)$ constructs $(G \cap C)$, that is, $G \cap C \in L(G \cap D)$.

Recall that when Ψ is an assertion in B 's forcing language (with L our ground model), Ψ 's “boolean value” or “ B -value”, denoted $\|\Psi\|_B$ (or just $\|\Psi\|$ when B is understood from context) is the (unique) element of B that will be in G (for any generic G) if and only if Ψ is true in $L[G]$.

3.2 Main Facts About Relative Constructibility

Constructibility, relative and otherwise, is reviewed in more detail in the appendix, Section 11. The present section states some fundamental results, referring to that appendix for the proofs, and then derives from these results some basic facts about the interconstructibility of reals, continua, and generic filters.

Definitions. A *standard* ZF model is one in which all elements are pure sets and “ \in ” is interpreted as the real membership relation among its elements. An *inner* ZF model is one that is transitive and has all the ordinals (we will allow “all the ordinals” to pass as an unproblematic concept prior to the appendix, Section 12). When y is any set, a *y -predicate* is a proposition $\Psi(x, c_1, \dots, c_n)$ of the language of set theory, having one free variable x and finitely many constant symbols c_1, \dots, c_n , considered together with fixed referents for the c_i chosen out of $L \cup \text{TrCl}(\{y\})$. A y -predicate $\Psi(x, c_1, \dots, c_n)$ is *y -absolute* if

$$\{x : \Psi(x, c_1, \dots, c_n)\}$$

evaluates to the same set in any standard inner ZF model having y as a member.

Lemma 3.1 *There is a proposition Ψ such that $\Psi(x, \alpha, y)$ is, for any ordinal α and set y , a y -absolute predicate satisfying $\{x : \Psi(x, \alpha, y)\} = L_\alpha(y)$.*

The proof is discussed in the appendix (Section 11). \square

Lemma 3.2 *$L(y)$ is the smallest standard inner ZF model having y as a member, in the sense that $L(y) \subseteq M$ for any other such model M .*

That $L(y)$ is a standard inner ZF model follows from a minor modification of the usual proof that L is such a model (see [6], Theorem 13.3, or [3], Theorem 1.2). The minimality claim for $L(y)$ follows from Lemma 3.1: If $L(y) \not\subseteq M$ then $L_\alpha(y) \not\subseteq M$ for some α , contradicting $L_\alpha(y)$'s definability by a y -absolute predicate. \square

Corollary 3.3 *The following are all easy consequences of Lemma 3.2:*

- (a) *If x constructs y (that is, $y \in L(x)$), then $L(y) \subseteq L(x)$.*
- (b) *The relation “ x constructs y ” ($y \in L(x)$) is transitive.*
- (c) *If x and y are interconstructible, then $L(x) = L(y)$ and $\mathbb{R}(x) = \mathbb{R}(y)$.*
- (d) *The ordering \leq_G on ACSAs(B) defined above (namely $C \leq_G D$ if and only if $G \cap C \in L(G \cap D)$) is guaranteed with boolean value 1 to be transitive; since it is also reflexive it is a preorder, so we may define $=_G, <_G$, etc. accordingly. \square*

It is a central fact about $L(y)$ that each of its members (not only each level $L_\alpha(y)$) is definable by a y -absolute predicate:

Lemma 3.4 $z \in L(y)$ if and only if $z = \{x : \Psi(x, c_1, \dots, c_n)\}$ for some y -absolute predicate $\Psi(x, c_1, \dots, c_n)$.

The proof is discussed in the appendix (Section 11.3). \square

We now bring these central facts about relative constructibility to bear on $L(G)$.

Lemma 3.5 For all (constructible) $X \subseteq B$, $G \cap \overline{X} \in L(G \cap X)$.

By Lemma 3.4, $G \cap \overline{X} \in L(G \cap X)$ if and only if $G \cap \overline{X}$ is definable by a $(G \cap X)$ -absolute predicate, which is what we will show.

\overline{X} , the boolean completion of X in B , was defined above as the smallest complete subalgebra of B containing X . The key to our proof is an alternative definition of \overline{X} as the union of a transfinite hierarchy.

When $Y \subseteq B$, define $\phi(Y)$ to be Y 's closure under arbitrary negations and joins in B , or more precisely:

$$\phi(Y) \equiv \{\neg z : z \in Y\} \cup \{\bigvee S : S \subseteq Y \text{ and } S \in L\}.$$

Now define a hierarchy by successive closures under ϕ :

$$X_0 \equiv X;$$

$$X_{\alpha+1} \equiv \phi(X_\alpha);$$

$$X_\alpha \equiv \bigcup_{\beta < \alpha} X_\beta \text{ when } \alpha \text{ is a limit ordinal.}$$

It is easily checked that the union of all the X_α is the smallest complete subalgebra of B that includes X . In fact the growth of the X_α 's must cease at some ordinal δ , lest \overline{X} 's cardinality exceed B 's, so we may say that \overline{X} is the union of the X_α 's with α less than this δ . The function $\Xi : \alpha \mapsto X_\alpha$, with domain $\delta + 1$, is a member of L . (Lemmas 11.2 and 11.4 can be used to establish $\Xi \in L(\emptyset)$ rigorously.) Note that $\Xi(0) = X$ and $\Xi(\delta) = \overline{X}$.

Now for all $\alpha \leq \delta$ define $\Gamma(\alpha) \equiv G \cap \Xi(\alpha)$. Plainly we have $\Gamma(0) = G \cap X$ and $\Gamma(\delta) = G \cap \overline{X}$. Because G is a generic ultrafilter on B , we will have for all $\alpha < \delta$:

$$\Gamma(\alpha + 1) = \{b \in \Xi(\alpha + 1) : \neg b \in \Xi(\alpha) \setminus \Gamma(\alpha), \text{ or } (\exists b' \in \Gamma(\alpha))(b' \leq b)\}.$$

(We have used here the fact that if $b = \bigvee S$ for some $S \in L, S \subseteq \Xi(\alpha)$, then those elements of B^+ that are \leq some S -member form a subset of B^+ that is dense below b ; hence by G 's genericity, if $b \in G$, some S -member must be in $\Gamma(\alpha)$.)

Note now that $\Gamma(\alpha)$ is the union of previous $\Gamma(\beta)$'s when α is a limit ordinal; and that because the expression for $\Gamma(\alpha + 1)$ above does not involve (either implicitly or explicitly) any unconstructible sets other than $\Gamma(\alpha)$, there is a proposition Ψ such that for all α , $\Psi(x, \Gamma(\alpha), \Xi, \alpha)$ is a $\Gamma(\alpha)$ -absolute predicate that defines $\Gamma(\alpha + 1)$. Lemmas 11.2 and 11.4 then ensure that $\Gamma(\delta)$ can be defined by a $\Gamma(0)$ -absolute predicate, and since $\Gamma(\delta) = G \cap \overline{X}$ and $\Gamma(0) = G \cap X$, we are done. \square .

Lemma 3.6 *Every real number $x \in L(G) \setminus L$ is interconstructible with $G \cap D$ for some $D \in \text{ACSAs}(B)$, and vice-versa.*

Fix $D \in \text{ACSAs}(B)$ and (since the definition of $\text{ACSAs}(B)$ requires D to be countably completely generated) fix $\{d_n : n \in \omega\} \subseteq D$ that completely generates D . By Lemma 3.5, $G \cap D \in L(G \cap \{d_n : n \in \omega\})$. If we define the real number x by $\{n : d_n \in G\}$, then x is clearly interconstructible with $G \cap \{d_n : n \in \omega\}$, so by Corollary 3.3 we have $G \cap D \in L(x)$. (And clearly $x \in L(G \cap D)$ since $x = \{n : d_n \in G \cap D\}$.)

Conversely, fix an unconstructible real number $x \subseteq \omega$. Let \dot{x} be a B -name for x such that $1 \Vdash \text{“}\dot{x} \subseteq \omega \text{ is unconstructible”}$, and let D be the subalgebra of B completely generated by $\{\|n \in \dot{x}\| : n \in \omega\}$. D will be atomless because an atom of D would force for all n either $n \in \dot{x}$ or $n \notin \dot{x}$, thus forcing x to be a member of the ground model L , contradicting the guarantee that \dot{x} names an unconstructible real. As in the previous paragraph, x will be interconstructible with $D \cap G$. \square

Lemma 3.7 *A real number x is interconstructible with its $\mathbb{R}(x)$.*

This is clear since $x \in \mathbb{R}(x) \subseteq L(\mathbb{R}(x))$, and $\mathbb{R}(x) \in L(x)$ by definition. \square

Lemma 3.8 *The singly-generated continua in a forcing extension $L[G]$ are precisely the sets of form $\mathbb{R}(G \cap D)$, where $D \in \text{ACSAs}(B)$ or $D = \emptyset$.*

$\mathbb{R}(\emptyset)$ is obviously singly-generated since we are construing real numbers as subsets of ω and \emptyset is such a subset.

Fix $D \in \text{ACSAs}(B)$. By Lemma 3.6, $D \cap G$ is interconstructible with some $x \subseteq \omega$; and by Corollary 3.3 (c), $\mathbb{R}(G \cap D)$ equals $\mathbb{R}(x)$, which is a singly-generated continuum. Conversely, fix an unconstructible singly-generated continuum $X \in L[G]$, and real number $x \subseteq \omega$ such that $X = \mathbb{R}(x)$. By the other direction of Lemma 3.6, x is interconstructible with some $G \cap D$, so again we will have $\mathbb{R}(x) = \mathbb{R}(G \cap D)$. \square

Corollary 3.9 *G is interconstructible with $\mathbb{R}(G)$ (since we assume B is countably completely generated).*

Our assumption that B is itself countably-completely-generated entails $B \in \text{ACSAs}(B)$, so by Lemma 3.6 G is interconstructible with some real number x . By Lemma 3.7, x is interconstructible with $\mathbb{R}(x)$. By transitivity, G is interconstructible with $\mathbb{R}(x)$. By Corollary 3.3, $\mathbb{R}(G) = \mathbb{R}(\mathbb{R}(x)) = \mathbb{R}(x)$. The claim follows. \square

Corollary 3.10 (General form of candidate families) *Stipulation 2.1 is equivalent to the assertion that every family of continua we will consider as a candidate to self-construct will have form*

$$\mathcal{F}(B, \zeta, G) \equiv \{\mathbb{R}(\emptyset)\} \cup \{\mathbb{R}(G \cap C) : C \in \text{ACSAs}(B) \text{ and } \zeta(C) \in G\},$$

where ζ is some mapping $\zeta : \text{ACSAs}(B) \rightarrow B$. \square

3.3 The Main Question

Plenty of forcing extensions $L[G]$ (including Cohen-forcing extensions) have instances of \mathcal{N} meeting requirements (i)-(iv) in the definition of self-collection into $\mathbb{R}(G)$. It is requirement (v), $\mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(G)$, that is hard to satisfy; in order to focus on it we assume the following:

Stipulation 3.11 *\mathcal{N} is a set of singly-generated continua meeting requirements (i)-(iv) in the definition of self-collecting into $\mathbb{R}(G)$, as well as Stipulation 2.1; furthermore, assume we have a name \dot{N} forced by $1 \in B$ to meet the foregoing.*

The question posed at the end of Section 2 now takes this form: Under what conditions will \mathcal{N} meeting Stipulation 3.11 satisfy the final requirement for self-collection, $\mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(G)$? This final requirement is, in light of the results of Section 3.2, equivalent to the following more perspicuous one:

Lemma 3.12 $\mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(G) \iff G \in L(\bigcup \mathcal{N})$ (under Stip. 3.11).

Clause (iii) of Stipulation 3.11 is $\mathcal{N} \in L(\mathbb{R}(G))$. By Corollary 3.9 and Corollary 3.3 (c), $L(G) = L(\mathbb{R}(G))$. Thus $\mathcal{N} \in L(G)$, and since $L(G)$ satisfies the ZF axiom of union, $\bigcup \mathcal{N} \in L(G)$. If also $G \in L(\bigcup \mathcal{N})$, then G is interconstructible with $\bigcup \mathcal{N}$, and by Corollary 3.3 (c) we have $\mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(G)$.

Conversely if $\mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(G)$ then $\mathbb{R}(G) \in L(\bigcup \mathcal{N})$, whence by Corollary 3.9 and transitivity (Corollary 3.3 (b)) $G \in L(\bigcup \mathcal{N})$. \square

This equivalence allows us to officially register our Main Question thus:

The Main Question (\mathcal{N} Version, Forcing-Extension Perspective): *Assuming Stipulation 3.11, under what conditions does $G \in L(\bigcup \mathcal{N})$ hold (so that \mathcal{N} fulfills the final requirement (v) for self-collection)?*

We will soon (Section 4) introduce a set Θ that is interconstructible with and more manageable than $\bigcup \mathcal{N}$, and we will find requirements on Θ equivalent to those of Stipulation 3.11 on \mathcal{N} . For the sake of keeping all versions of our Main Question in one place, we will register the second one now.

The Main Question (Θ Version, Forcing-Extension Perspective): *Assuming Stipulation 3.11, under what conditions does $G \in L(\Theta)$ hold?*

We will answer this version of the question in Theorem 7.8.

Note now that the question “Under what conditions does \mathcal{N} self-collect into $\mathbb{R}(G)$?” is posed from the perspective of the forcing extension $L[G]$, and that we would ultimately

like to answer a question posed from the perspective of the ground model L : “Under what conditions on B and on the B -name $\dot{\mathcal{N}}$ will $\dot{\mathcal{N}}$ ’s interpretation be *forced* to self-collect into $\mathbb{R}(G)$?”

The Main Question (Ground-Model Perspective): *Assuming Stipulation 3.11, how do we calculate $\|\dot{G} \in L(\dot{\mathcal{N}})\|_B$?*

We will answer this version of the question in Theorem 8.7.

As to the question of whether these conditions can ever be met, the answer is yes; we will present in Section 10 an example of a set \mathcal{N} of continua that self-collects. However, this example does not provide a model of the self-construction axioms since for each $X \in \mathcal{N}$, no subset of X ’s predecessors (that is, of X ’s proper sub-continua) self-collects into X . Prospects for obtaining models of the self-construction axioms are discussed in Section 10.

4 Equivalence Classes $\theta(X)$ Of Generic Filters

We now present the key to our analysis, which is a shift in perspective from reals and sets of reals, to sets of generic filters with which they are interconstructible.

Our Main Question (in its first version above) is how to arrange for the generic filter G to be constructible from the set $\dot{\mathcal{N}}$ of reals. We know from Lemma 3.6 that each real number $x \in \dot{\mathcal{N}}$ will be interconstructible with the filter $G \cap D$ for some subalgebra D of B . This suggests the following.

The naïve idea: $G \in L(\dot{\mathcal{N}})$ will hold if and only if there is some absolute way to define a function mapping each $x \in \dot{\mathcal{N}}$ to “the” filter $G \cap D$ that is interconstructible with x , such that the union of all these filters is G .

The problem with this idea, indicated by the scare-quotes around “the,” is that multiple generic filters F on subalgebras of B might be interconstructible with a real x , not all of them having form $G \cap D$. In general, the closest we can get to realizing the hope just expressed is to associate each singly-generated continuum $Y \in L(\dot{\mathcal{N}})$ with an *equivalence class* of generic filters interconstructible with Y . We call this equivalence class Y ’s “constructibility degree” of filters, and denote it $\theta(Y)$.

Definitions.

$$\theta(Y) \equiv \{F \in L(Y) : F \text{ is a } L\text{-generic filter on some } C \in \text{ACSAs}(B) \text{ and } Y \in L(F)\}.$$

We now let Θ be an interconstructible stand-in for $\dot{\mathcal{N}}$ built with these constructibility degrees, defined in the following $\dot{\mathcal{N}}$ -absolute way:

$$\Theta \equiv \{\theta(\mathbb{R}(x)) : x \in \dot{\mathcal{N}}\}.$$

Θ is indeed interconstructible with $\bigcup \mathcal{N}$, since in the other direction we have

$$\bigcup \mathcal{N} = \bigcup \{\mathbb{R}(M) : M \in \Theta\}.$$

It follows from the transitivity of relative constructibility (Corollary 3.3) that a given set is constructible from Θ if and only if it is constructible from $\bigcup \mathcal{N}$, and, in particular:

Lemma 4.1 *Under Stipulation 3.11, \mathcal{N} satisfies $G \in L(\bigcup \mathcal{N})$ if and only if $G \in L(\Theta)$, where Θ is defined in terms of $\bigcup \mathcal{N}$ as above. \square*

This confirms the equivalence of the Main Question’s first and second versions. Our hope for answering the latter — i.e. for finding a condition under which G is constructible from Θ — is a less naïve version of the idea expressed above.

The right idea: $G \in L(\Theta)$ will hold just if there is something about Θ ’s structure that lets us deduce (using a single filter $G \cap D \in \bigcup \Theta$ as a “hint”, if necessary) which filters in $\bigcup \Theta$ are genuinely of form $G \cap D$, as opposed to specious “copies,” and lets us conclude that G is the union of all the former.

Essentially all of our work from this point through Theorem 7.8 is dedicated to realizing this idea. We stress this now, lest it be obscured by the technical details leading up to that Theorem.

Since \mathcal{N} meets Stipulation 3.11 with boolean (B) value 1, it is easy to verify that Θ as we have defined it from \mathcal{N} will meet the following with boolean value 1:

Lemma 4.2 *Under Stipulation 3.11, the following assertions, corresponding to the first three requirements of the Self-Collection axiom, are forced by $1 \in B$:*

- (i). $G \notin \bigcup \Theta$;
- (ii). $(G \cap C, G \cap D \in \bigcup \Theta) \iff G \cap (\overline{C \cup D}) \in \bigcup \Theta$;
- (iii). $\Theta \in L(G)$ [trivial since Θ is G ’s interpretation of a B -name]. \square

Structure of the Equivalence Classes $\theta(X)$

We now show that generic filters on ACSAs(B)-members in $L[G]$ are interconstructible if and only if they are “essentially isomorphic copies” of each other; consequently, the constructibility degrees $\theta(Y)$ will be equivalence classes of copies. (We should note that the basic idea here has long been understood; we introduce the term “copy” only as a convenience.)

Definitions:

$a, b \in B$ are *compatible* if $a \wedge b \neq 0$; similarly, a is compatible with $X \subseteq B$ if it is compatible with each X -member, and $X, Y \subseteq B$ are compatible if they are pairwise compatible.

The *upwards closure* of $X \subseteq B$ is $\{b \in B : (\exists x \in X)(x \leq b)\}$.

A *principal ideal* of $C \in \text{ACSAs}(B)$ is $\{c \in C : c \leq e\}$ for some $e \in C^+$ (we put C^+ rather than C here because it will simplify matters to exclude the degenerate ideal $\{0\}$ by definition).

When $c \in C^+$, we write $C \upharpoonright c$ to mean the principal ideal of C whose greatest member is c ; and we write “ $\max(I)$ ” for the greatest member of a principal ideal I .

Similarly, when ϕ is a function on C , we write $\phi \upharpoonright c$ to mean ϕ 's restriction to $C \upharpoonright c$. (When context suggests that X is not a member of C , but rather a set likely to have a nonempty intersection with C , $\phi \upharpoonright X$ will mean ϕ 's restriction to $C \cap X$, as usual.)

Note that a principal ideal $C \upharpoonright c$ can be considered a boolean algebra in its own right, having c as its greatest member, inheriting \wedge and \vee from C , and having a negation operation defined from C 's negation \neg as $e \mapsto c \wedge \neg e$.

When ϕ is a function, $\text{dom } \phi$ and $\text{ran } \phi$ denote its domain and range; and we adopt the common square-bracket notation $\phi[X]$ to mean $\text{ran}(\phi \upharpoonright X)$.

If $b \in B$ and $A \subseteq B$, then $b \wedge [A]$ means $\{b \wedge a : a \in A\}$.

$\text{Aut}(B)$ is the set of automorphisms of B (bijections from B onto itself that preserve boolean structure) in the ground model L .

A *partial automorphism* of B is a bijection between principal ideals of (possibly distinct) ACSAs of B that preserves boolean structure (when the principal ideals are considered as boolean algebras in their own right, as described above).

$\text{ParAut}(B)$ is the set of B 's partial automorphisms in L .

Definition of “copy”: When $C, D \in \text{ACSAs}(B)$, and F is a filter on C , and $\phi \in \text{ParAut}(B)$ is an isomorphism from some principal ideal of C not disjoint from F to some principal ideal of D , we say that ϕ *copies* F to the upward closure in D of $\phi[F \cap \text{dom } \phi]$, and that this upward closure is a *copy* of F , *via* ϕ .

The importance of copies is established in the following lemma, which shows that for any real number x , its “constructibility degree” $\theta(\mathbb{R}(x))$ of filters is just an equivalence class of copies.

Lemma 4.3 *For each singly-generated unconstructible continuum $X \in L[G]$,*

$(\exists C)(G \cap C \in \theta(X))$, and

$(\forall C)(G \cap C \in \theta(X) \Rightarrow \theta(X) = \{F : F \text{ is a copy of } G \cap C\})$.

We will establish some preliminary lemmas in order to prove this.

Lemma 4.4 *If G_C, G_D are generic filters on (respectively) $C, D \in \text{ACSAs}(B)$, then G_C constructs G_D (i.e. $G_D \in L(G_C)$) if and only if there exists $C' \in \text{ACSAs}(B)$, $C' \subseteq C$, such that G_D is a copy of $G_C \cap C'$.*

The “if” direction is straightforward; we consider the “only if” direction. Suppose $G_D \in L(G_C)$. G_D is interconstructible with a real number r_D by Lemma 3.6. Therefore by transitivity of relative constructibility (Corollary 3.3 (b)), $r_D \in L(G_C)$. Again by Lemma 3.6, this time using our C for its B and our C_G for its G , we have r_D interconstructible with $L(G_C \cap C')$ for some $C' \in \text{ACSAs}(C)$. Then by Corollary 3.3 (b) and (c), G_D is interconstructible with $G_C \cap C'$, and $L(G_D) = L(G_C \cap C')$.

Note that G_D and $(G_C \cap C')$ are subsets of L , so as we have mentioned the parentheses and square-brackets versions of relative constructibility are equivalent for them, i.e. $L(G_D) = L[G_D]$ and $L(G_C \cap C') = L[G_C \cap C']$.

Since $G_D \in L[G_C \cap C']$, there exists a C' -name γ whose interpretation by $G_C \cap C'$ is G_D . Being a filter, G_D is upwards-closed in D , and therefore the following “nice” C' -name δ also yields G_D when interpreted by $G_C \cap C'$:

$$\delta \equiv \{(\check{d}, \bigvee \{\|\check{d}' \in \gamma\|_{C'} : d' \in D \text{ and } d' \leq d\}) : d \in D\}.$$

We will use δ rather than γ as a name for G_D because δ , considered as a function whose domain is the set of canonical forcing names for elements of D , satisfies $d \leq d' \Rightarrow \delta(\check{d}) \leq \delta(\check{d}')$. Similarly, let σ be a nice D -name whose interpretation by G_D is $G_C \cap C'$, and which satisfies $c \leq c' \Rightarrow \sigma(\check{c}) \leq \sigma(\check{c}')$.

For the remainder of this proof we will, for simplicity, conflate elements of D with their canonical C' -names, and vice-versa, so that δ can be considered a function from D into C' , and σ a function from C' into D . Therefore we may assert the following:

$$d \leq d' \Rightarrow \delta(d) \leq \delta(d'); \tag{1}$$

$$c \leq c' \Rightarrow \sigma(c) \leq \sigma(c'); \tag{2}$$

$$d \in G_D \iff \delta(d) \in G_C \cap C'; \tag{3}$$

$$c \in G_C \cap C' \iff \sigma(c) \in G_D. \tag{4}$$

From the last two properties we have

$$(\forall c \in C')(c \in G_C \iff \delta(\sigma(c)) \in G_C). \tag{5}$$

Now the genericity of G_C would guarantee a counterexample to (5) unless there existed $c \in G_C \cap C'$ such that:

$$\delta(\sigma(c')) = c' \text{ for all } c' \in C' \upharpoonright c. \tag{6}$$

Similarly, because G_D is generic, we deduce the existence of $d \in G_D$ such that:

$$\sigma(\delta(d')) = d' \text{ for all } d' \in D \upharpoonright d. \tag{7}$$

Since $d \in G_D$ we have $\delta(d) \in G_C$ so that $\delta(d) \wedge c \in G_C \cap C'$.

We claim that σ 's restriction to $C' \upharpoonright (\delta(d) \wedge c)$ is a $\text{ParAut}(B)$ -member that copies $(G_C \cap C')$ to G_D . It is an isomorphism, that is, $c_1 \leq c_2 \iff \sigma(c_1) \leq \sigma(c_2)$, for (2) above

establishes one direction, and in the other, if $c_1 \not\leq c_2$ but $\sigma(c_1) \leq \sigma(c_2)$, then (6) and (1) entail a contradiction. Finally, the range of $\sigma \upharpoonright \delta(d) \wedge c$ is all of $D \upharpoonright \sigma(\delta(d) \wedge c)$; if $d' < d$ were a counterexample to this, $\delta(d')$ would be in the domain of $\sigma \upharpoonright \delta(d) \wedge c$ by (1), but then $\sigma(\delta(d')) \neq d'$, violating (7). \square

Lemma 4.5 *If in addition to the supposition of Lemma 4.4 we have $L(G_D) = L(G_C)$, we can require there that C' be C itself.*

At the outset of our proof we chose C' to be any ACSA of C such that $(G_C \cap C')$ is interconstructible with G_D , so knowing $L(G_D) = L(G_C)$ allows us to choose C itself. \square

Lemma 4.6 *A copy F' of a filter F on some $C \in \text{ACSAs}(B)$ will be L -generic if and only if F is L -generic.*

Let ϕ copy F to F' , and let $D \in \text{ACSAs}(B)$ be such that $\text{ran } \phi$ is a principal ideal of D , and F' is the upwards closure of $\phi[F \cap \text{dom } \phi]$ in D .

We first show that F will be L -generic on C —that is, meet every constructible dense subset S of C^+ —just if $F \cap \text{dom } \phi$ is L -generic on $\text{dom } \phi$. This is trivial if $\max(\text{dom } \phi) = 1$, so assume otherwise. Note that $\max(\text{dom } \phi) \in F$ by the “ F not disjoint from $\text{dom } \phi$ ” clause of the definition of “copy,” and the fact that filters are upwards-closed.

Suppose F is L -generic on C and let S be any constructible dense subset of $\text{dom } \phi$. Define $S' \equiv S \cup (C^+ \upharpoonright \neg \max(\text{dom } \phi))$. S' is dense in C^+ , so there exists $s \in F \cap S'$; and this s must be in the S half of S' since otherwise, by the closure of F under \wedge , we would have $\max(\text{dom } \phi) \wedge s = 0 \in F$, contradicting F 's being a proper filter. Conversely, if $F \cap \text{dom } \phi$ is L -generic on $\text{dom } \phi$, and S is any constructible dense subset of C^+ , define $S' \equiv \max(\text{dom } \phi) \wedge [S]$; then S' is dense in $\text{dom } \phi$, so there exists $s' \in F \cap S'$, which by definition of S' must be of form $s' = \max(\text{dom } \phi) \wedge s$ for some $s \in S$, and this $s \in S$ must also be in F by the upward-closure of filters.

Likewise, F' is L -generic on D just if $F' \cap \text{ran } \phi$ is L -generic on $\text{ran } \phi$. The lemma then follows from the fact that ϕ is an isomorphism, which entails that a subset of $\text{dom } \phi$ will be dense just if its image is dense in $\text{ran } \phi$. \square

Proof of Lemma 4.3:

“ $\exists C$ ” clause: Just after defining $\theta(X)$ above, we noted that it always has a member of form $G \cap C$ (unless X is constructible, a case that Lemma 4.3's statement explicitly excludes).

“ $\forall C$ ” clause: Letting C be arbitrary such that $G \cap C \in \theta(X)$, $G \cap C$ is of the form stated in the lemma because it is trivially a copy of itself; any other copy of $G \cap C$ is L -generic by Lemma 4.6 and is clearly interconstructible with $G \cap C$, so satisfies the definition of a

$\theta(X)$ -member. Finally, it follows from Lemma 4.4 and Lemma 4.5 that every $F \in \theta(X)$ is of this form. \square

The following lemma gives more information about an important special case of Lemma 4.4, in which G_C and G_D are the restrictions to C and D of our generic G on B .

Lemma 4.7 $D \leq_G C \iff (\exists g \in G)(g \wedge [D] \subseteq g \wedge [C])$.

Recall that $D \leq_G C$ by definition means $G \cap D \in L(G \cap C)$, and that, because $G \cap C \subseteq L$, we have $L(G \cap C) = L[G \cap C]$.

\Rightarrow : We are given $G \cap D \in L[G \cap C]$, so there is a C -name (which we can regard as a B -name involving only elements of C) whose interpretation by $G \cap C$ is $G \cap D$; call this name δ . Because δ names the same set as the canonical B -name $\dot{G} \cap \check{D}$, we must have

$$\|\delta = \dot{G} \cap \check{D}\|_B \in G.$$

Let g denote $\|\delta = \dot{G} \cap \check{D}\|_B$; we will verify that for all $d \in D$ some $c \in C$ satisfies $c \wedge g = d \wedge g$. We claim in fact that for all d , $\|\check{d} \in \delta\|_C$ is a c that satisfies this. For if $\|\check{d} \in \delta\|_C \wedge g \neq d \wedge g$, then

$$g \wedge ((d \wedge \neg \|\check{d} \in \delta\|_C) \vee (\neg d \wedge \|\check{d} \in \delta\|_C))$$

is nonzero, and is an element of B below g that forces $\delta \neq \dot{G} \cap \check{D}$, impossible.

\Leftarrow : Consider arbitrary $d \in D$; it is given that some $c \in C$ satisfies $c \wedge g = d \wedge g$. If some such c is in $G \cap C$, then $g \wedge c \in G$, and $g \wedge c \leq d$, so $D \in G$. Otherwise some such c is not in $G \cap C$, which entails $c \wedge g \notin G$, which entails $d \notin G$. Therefore $\{d \in D : (\exists c \in G \cap C)(g \wedge c = g \wedge d)\}$ will evaluate to $G \cap D$ in $L(G \cap C)$. \square

We can express the meaning of Lemma 4.7 by saying: there is some element g of information in the filter G , such that any *additional* information about G 's extension that might be supplied by $G \cap D$, could just as well be supplied by $G \cap C$. This suggests the following terminology.

Definition: Call $b \in B^+$ an *information supplement for C 's generic to construct D 's generic* if $b \wedge [D] \subseteq b \wedge [C]$. Let $IS(C, D)$ denote the set of all such information supplements; it is clearly a downward-closed subset of B^+ . With this definition we can restate Lemma 4.7 thus:

Corollary 4.8 $\|C \geq_G D\|_B = \bigvee IS(C, D)$. \square

5 Self-Collection Condition In Terms Of Definability

We are now ready to begin answering the Main Question (in its “ Θ version”) stated at the end of Section 3.3, from the perspective of the generic extension $L[G]$. The question is under what conditions Θ obeying Stipulation 3.11 (and therefore all clauses of Lemma 4.2) constructs G , meaning $G \in L(\Theta)$.

Our starting point is an answer we have already found, in terms of definability: By Lemma 3.4, $G \in L(\Theta)$ if and only if G is definable by a Θ -absolute predicate. This answer is not very helpful by itself, since there is no obvious way to check whether this condition holds. We will now begin the work of translating it into a more easily verifiable condition, work which will culminate in Theorem 7.8. The first step is to obtain a somewhat tighter definability condition.

Lemma 5.1 *Under Stipulation 3.11, $G \in L(\Theta)$ if and only if*

$$G = \{x : \Psi(x, \Theta, G \cap C, c'_1, \dots, c'_n)\}$$

for some Θ -absolute predicate $\Psi(x, \Theta, G \cap C, c'_1, \dots, c'_n)$ such that each c'_i is a constant in L and $G \cap C \in \bigcup \Theta$.

The “only if” direction is the nontrivial one. Assume $G \in L(\Theta)$ and (invoking Lemma 3.4) that G is definable by a Θ -absolute predicate as

$$G = \{x : \Delta(x, c_1, \dots, c_n)\}.$$

We wish to show that there exists a Θ -absolute predicate $\Psi(\dots)$ of the more narrowly-defined form demanded in the lemma’s statement, that defines the same set G .

The idea here is as follows. Δ ’s given constants c_1, \dots, c_n must refer, by the definition of Θ -absolute predicate, to sets that belong to L and/or to $\text{TrCl}(\{\Theta\})$. We wish to find a *single* constant referring to some $(G \cap C) \in \text{TrCl}(\{\Theta\})$, such that each c_i can be replaced in Δ by a subformula that defines c_i ’s referent using only Θ , this $G \cap C$, and finitely many constants *required to be members of L* , all of which will be parameters for our Ψ . What allows us to do this is the fact (established by Lemma 4.3) that each given c_i that is not already Θ or an L -member must be interconstructible with $(G \cap C_i)$ for some $C_i \in \text{ACSAs}(B)$ such that $(G \cap C_i) \in \bigcup \Theta$. Clause (ii) of Lemma 4.2 on Θ (which translates the directedness requirement on \mathcal{N}) then guarantees the existence of a single $C \in \text{ACSAs}(B)$ that includes each such C_i , and satisfies $(G \cap C) \in \bigcup \Theta$. Each of the $(G \cap C_i)$ ’s is then definable as the intersection with C_i of this one $(G \cap C)$. And each C_i is an L -member and so can be appended as another constant for our Ψ .

We formalize this argument in the appendix, Section 11. \square

A key consequence of Lemma 5.1 is that the predicate’s Θ -absoluteness rules out the existence of certain filters G' whose restriction to a certain C is the same as G ’s. This is made precise in the following lemma.

Lemma 5.2 *If the situation stated in Lemma 5.1 holds, and is forced to hold (as it must be) by some $g \in G$, i.e.*

$$g \Vdash \dot{G} = \{\check{x} \in \check{B} : \Psi(\check{x}, \dot{\Theta}, \dot{G} \cap \check{C}, \check{c}'_1, \dots, \check{c}'_n)\}, \quad (\star)$$

then there cannot exist in any ZF model an L -generic filter G' on B , $G' \neq G$, such that $g \in G'$, $G' \cap C = G \cap C$, and G' gives the same interpretation as G to the B -name $\dot{\Theta}$.

For if such a G' existed, and were used to interpret B -names to yield a generic extension $L[G']$, then the expression

$$\{x \in B : \Psi(x, \Theta, G \cap C, c'_1, \dots, c'_n)\}$$

would evaluate there to $G' \neq G$, contradicting the Θ -absoluteness of $\Psi(\dots)$. \square

6 Definition of χ and rigid inclusions of boolean algebras

We have just seen in Lemma 5.2 that if $G \in L(\Theta)$ is to hold, there cannot exist certain filters $G' \neq G$ having the same restriction to some subalgebra C as G . We will now see that, in the special case that the g in that lemma equals 1, this also rules out certain automorphisms of B that leave C fixed and would map G onto such a forbidden G' . This suggests the idea of a *rigid inclusion* of boolean algebras $C \subseteq B$, for which no such automorphisms exist. The fact we will prove in this section makes this idea precise. Because this fact concerns only the special case just noted (i.e. $g = 1$) it will not be referenced later. We include it nonetheless because it shows in a particularly simple way the relevance to our work of rigid inclusions, whose importance will become clearer later: in Section 9.1 we will see that a lack of rigid boolean inclusions $C \subset B$, combined with a few other properties, suffices to prevent self-collection; and in Section 10 we will propose a way to obtain rigid boolean inclusions, as the most promising route to a model of the self-construction axioms.

Definition. We begin by defining a mapping $\chi : \text{ACSAs}(B) \rightarrow B$ that will be central to the remainder of this paper; χ encodes the conditions under which, for each D , $G \cap D$ will be a member of $\bigcup \Theta$:

$$\chi(D) \equiv \|\dot{G} \cap D \in \bigcup \dot{\Theta}\|_B.$$

Lemma 6.1 *Under Stipulation 3.11 the following hold:*

- (i). $\chi(B) = 0$;
- (ii). $(\forall C, D)(\chi(C) \wedge \chi(D) = \chi(\overline{C \cup D}))$.

This is easy to verify since each claim about χ is a straightforward translation of the corresponding one on Θ in Lemma 4.2. \square

Lemma 6.2 (i) $(G \cap C \in \bigcup \Theta \text{ and } C \geq_G D) \Rightarrow G \cap D \in \bigcup \Theta;$

(ii) $(\forall C, D)(\chi(C) \wedge \|C \geq_G D\| \leq \chi(D)).$

For (i), suppose $G \cap C \in \bigcup \Theta$ and $C \geq_G D$. By Lemma 3.6, let y be a real number interconstructible with $G \cap D$. By the definition of Θ , a generic filter will be a member of $\bigcup \Theta$ if and only if it is interconstructible with some real number x in some continuum $Z \in \mathcal{N}$; fix witnesses x, Z for $G \cap C$. By the definition of \geq_G and the transitivity of relative constructibility (Lemma 3.3), x constructs y ; thus $y \in Z$. Then y, Z witness $G \cap D \in \bigcup \Theta$.

Claim (ii) follows easily from (i) and the definition of χ . \square

Definitions. $C \subseteq B$ is a *rigid boolean inclusion* if no $\phi \in \text{Aut}(B)$ other than the identity mapping leaves every C -member fixed. It is a *rigid boolean+ χ inclusion* if no $\phi \in \text{Aut}(B)$ other than the identity mapping leaves every C -member fixed and preserves χ , in the sense that $\phi(\chi(D)) = \chi(\phi[D])$ for all $D \in \text{ACSAs}(B)$. (Note that if $C \subseteq B$ is a rigid boolean inclusion then it is also a rigid boolean+ χ inclusion, for any χ .)

Fact 6.3 Recall that (\star) of Lemma 5.2 must hold for some C and g if we are to have $G \in L(\Theta)$; if (\star) holds for some C and $g = 1$, then $C \subseteq B$ is a rigid boolean+ χ inclusion. If furthermore $\chi(D) = \|D <_G B\|$ for all $D \in \text{ACSAs}(B)$, then $C \subseteq B$ is a rigid boolean inclusion.

Suppose (\star) of Lemma 5.2 holds with $g = 1$ for some fixed C .

For the first claim, suppose $\phi \in \text{Aut}(B)$ furnished a counterexample, i.e. ϕ is not the identity mapping but preserves χ and leaves each C member fixed. Let G be a generic ultrafilter on B such that when we define $G' \equiv \phi[G]$, we have $G' \neq G$. (This is possible because ϕ is not the identity mapping.) Consider G' . It too is L -generic, by Lemma 4.6. Since ϕ leaves every C member fixed, we have $G \cap C = G' \cap C$. And because ϕ exists in the ground model and satisfies $\phi(\chi(D)) = \chi(\phi[D])$ for all $D \in \text{ACSAs}(B)$, $\dot{\Theta}$'s interpretation is the same under G' as under G , as is easily verified using χ 's definition in terms of $\dot{\Theta}$. But then since (\star) holds with $g = 1$, G' violates Lemma 5.2; so the first claim can have no counterexample.

For the second claim, suppose $\chi(D) = \|D <_G B\|$ for all $D \in \text{ACSAs}(B)$; it suffices to verify that this implies every $\phi \in \text{Aut}(B)$ preserves χ , for then any counterexample ϕ to $C \subseteq B$ being a rigid boolean inclusion would also be a counterexample to its being a rigid boolean+ χ inclusion.

Let $\phi \in \text{Aut}(B)$ and $D \in \text{ACSAs}(B)$ be arbitrary. From the definition of $IS(C, D)$ used in Corollary 4.8,

$$IS(D, B) = \{b \in B^+ : B \upharpoonright b = b \wedge [D]\}.$$

It follows from this and the fact that ϕ is a boolean isomorphism that for all b ,

$$b \in IS(D, B) \iff \phi(b) \in IS(\phi[D], B),$$

which in turn implies

$$\phi(\bigvee IS(D, B)) = \bigvee IS(\phi[D], B).$$

Then by Corollary 4.8 and the fact that $\|D \geq_G B\| = \neg\|D <_G B\|$,

$$\phi(\neg\|D <_G B\|) = \neg\|\phi[D] <_G B\|.$$

Since ϕ preserves \neg ,

$$\phi(\|D <_G B\|) = \|\phi[D] <_G B\|;$$

then our supposition that $\chi(D) = \|D <_G B\|$ and $\chi(\phi[D]) = \|\phi[D] <_G B\|$ implies

$$\phi(\chi(D)) = \chi(\phi[D]).$$

Since D was arbitrary, this means ϕ preserves χ , which, as we noted, is what it suffices to verify. \square

We do not know at present whether a *sufficient* condition for self-collection can be found in terms of the existence of rigid boolean+ χ inclusions. Matters would be greatly simplified if it were. In our proof of Theorem 7.8, though, we will see that it may be possible to cobble together some of B 's *partial* automorphisms—in a model larger than $L[G]$ —to obtain a filter G' as in Lemma 5.2 that witnesses the failure of self-collection, yet is not isomorphic to G via any ground-model automorphism of B .

In any case, the existence of a rigid boolean+ χ inclusion $C \subset B$ would not supply any obvious formula Ψ for defining G from Θ and $G \cap C$ (plus finitely many constants in L), as Lemma 5.1 requires. In the next sections we will find a necessary and sufficient condition for self-collection that does provide such a Ψ .

7 Self-Collection Condition in Terms of Filters

The concept we use for turning Lemma 5.1's self-collection condition into a practically verifiable one is “generic refinability.” We will define this concept in parallel for generic filters on ACSA's of B , and for $\text{ParAut}(B)$ -members. Having done so, we will derive a self-collection condition in terms of generically-refinable filters that can be checked from the generic-extension perspective; then in the next section we will translate it into a condition in terms of generically-refinable $\text{ParAut}(B)$ -members, which can be checked in the ground model perspective. A basic technical tool we will need is “one-element refinements.”

7.1 One-element refinements

We will often need to expand an ACSA by adding a single B -member to it and taking its completion. We use the following notation for this: When $C \in \text{ACSAs}(B)$ and $b \in B$,

$$C^{+b} \equiv \{(c_1 \wedge b) \vee (c_2 \wedge \neg b) : c_1, c_2 \in C\}.$$

Lemma 7.1 *If $C \in \text{ACSAs}(B)$ and $b \in B$, then $C^{+b} \in \text{ACSAs}(B)$.*

This follows from routine boolean-algebraic manipulations. \square

Definitions. If F, F' are filters, we say F' *refines* F if $F \subseteq F'$. If $\phi, \phi' \in \text{ParAut}(B)$, we say ϕ' *refines* ϕ if:

$$\max(\text{dom } \phi') \leq \max(\text{dom } \phi), \text{ and } \max(\text{ran } \phi') \leq \max(\text{ran } \phi); \text{ and}$$

$$(\forall e \in \text{dom } \phi)(\phi'(e \wedge \max(\text{dom } \phi')) = \phi(e) \wedge \max(\text{ran } \phi')).$$

Note the second clause implies $\max(\text{dom } \phi') \wedge [\text{dom } \phi] \subseteq \text{dom } \phi'$.

When $b \in B$ and C is an ACSA of B — or, more generally, a subset of B that is closed under \bigwedge (arbitrary meets) — we define

$$b \uparrow C \equiv \bigwedge \{c \in C : b \leq c\}.$$

Given $\phi \in \text{ParAut}(B)$ and some $b \in B^+$, $b \leq \max(\text{dom } \phi)$, a *one-element domain refinement* of ϕ by b is a refinement ϕ^{+b} of ϕ satisfying $\text{dom } \phi^{+b} = b \wedge [\text{dom } \phi]$ and $\max(\text{ran } \phi^{+b}) = \phi(b \uparrow \text{dom } \phi)$. We will now see that these requirements uniquely determine ϕ^{+b} .

Lemma 7.2 *If $C \in \text{ACSAs}(B)$ and $b \in B^+$, the function $\psi : a \mapsto a \wedge b$ with $\text{dom } \psi = C \uparrow (b \uparrow C)$ is an isomorphism such that $\psi \in \text{ParAut}(B)$, and $\psi(a) \uparrow C = a$ for all $a \in \text{dom } \psi$.*

If we can show

$$(a \wedge b) \uparrow C = a \tag{*}$$

for all $a \in \text{dom } \psi$, then the lemma's other claims will follow easily. Let $a \in \text{dom } \psi$ be arbitrary, so that $a \in C$ and $a \leq (b \uparrow C)$. Plainly (*) holds with \leq in place of $=$; we must show that this inequality cannot be strict. So suppose

$$(a \wedge b) \uparrow C = a' < a$$

holds for some a' . It would follow that $b \wedge a = b \wedge a'$. Then boolean laws entail

$$\begin{aligned} b &= (b \wedge a) \vee (b \wedge \neg a) \\ &= (b \wedge a') \vee (b \wedge \neg a) \\ &= b \wedge (a' \vee \neg a). \end{aligned}$$

Thus $b \leq (a' \vee \neg a)$. But then also $b \leq (a' \vee \neg a) \wedge (b \uparrow C)$; the right side of that inequality is strictly $< (b \uparrow C)$, contradicting the definition of $(b \uparrow C)$ as the least C -member that is $\geq b$. \square

Lemma 7.3 *Given $\phi \in \text{ParAut}(B)$ and some $b \in B^+$, $b \leq \max(\text{dom } \phi)$, the one-element domain refinement ϕ^{+b} exists and is unique.*

Let C be any ACSA of B such that $\text{dom } \phi \subseteq C$, and note that $b \wedge [\text{dom } \phi]$ ($= b \wedge [C]$) is a principal ideal of C^{+b} defined above.

By Lemma 7.2, the map $\psi : a \mapsto a \wedge b$ is an isomorphism mapping $C \upharpoonright (b \upharpoonright C)$ onto $b \wedge [C]$, the latter being a principal ideal of C^{+b} . Thus we may define ϕ^{+b} by composition:

$$\phi^{+b}(d) \equiv \phi(\psi^{-1}(d)),$$

with the domain of ϕ^{+b} set to $b \wedge [C]$. It is straightforward to check that ϕ^{+b} meets the definition of a refinement of ϕ , and that it is the unique such refinement satisfying the definition of a one-element refinement of ϕ by b . \square

Definition. In light of Lemma 7.3 we may use ϕ^{+b} to denote *the* one-element domain refinement of ϕ by b . Similarly, for $b \leq \max(\text{ran } \phi)$, we define the *one-element range refinement* ϕ_{+b} of ϕ to be the refinement of ϕ satisfying $\text{ran } \phi_{+b} = b \wedge [\text{ran } \phi]$ and $\max(\text{dom } \phi_{+b}) = \phi^{-1}(b \upharpoonright \text{ran } \phi)$.

Lemma 7.4 *For all $C \in \text{ACSAs}(B)$ and $b \in B^+$, $\|C =_G C^{+b}\| = 1$; so $\chi(C) = \chi(C^{+b})$.*

Suppose $b \in G$. By Lemma 7.2, the map $a \mapsto a \wedge b$ is a member of $\text{ParAut}(B)$ mapping $C \upharpoonright (b \upharpoonright C)$ onto $b \wedge [C]$, the latter being a principal ideal of C^{+b} . Since $b \in G$, neither the domain of this map nor its range is disjoint from G , so this map copies $G \cap C$ to $G \cap C^{+b}$, and its inverse copies $G \cap C^{+b}$ to $G \cap C$. Thus by Lemma 4.4, $G \cap C^{+b}$ and $G \cap C$ are interconstructible; whence by the definition of $=_G$ we have $C =_G C^{+b}$.

If on the other hand $\neg b \in G$, the same argument goes through with the map $a \mapsto a \wedge \neg b$. Thus both b and $\neg b$ force $C =_G C^{+b}$; thus 1 forces this too. $\chi(C) = \chi(C^{+b})$ then follows from Lemma 6.2. \square

7.2 Generic Refinability In Accordance With χ

The definitions of Γ_α , Γ , Φ_α , and Φ below are cumbersome, because they use recursion; it is a question we have not been able to settle whether there are simpler non-recursive definitions that would be equivalent. We will see that if $\Phi_1 = \Phi_0$, then $\Phi = \Phi_0$ (Lemma 9.4), but we have found no comparable simplification for other cases.

Accordance with χ . If F is an ultrafilter on $C \in \text{ACSAs}(B)$, we say it *accords with χ* if $\chi(C)$ is in the upwards closure of F in B . For $\phi \in \text{ParAut}(B)$, we say ϕ *accords with χ* if its domain and range are principal ideals of some C and D (respectively) such that $\max(\text{dom } \phi) \leq \chi(C)$ and $\max(\text{ran } \phi) \leq \chi(D)$. Note C and D may not be uniquely determined (since two distinct subalgebras of B can have a principal ideal in common), but the definition of χ ensures that if the requirement just stated holds for one choice of C, D , it holds for all choices. This is because whenever $\text{dom } \phi$ is a principal ideal of both C and C' , we have $\max(\text{dom } \phi) \Vdash C =_G C'$ (by Lemma 4.7), so by Lemma 6.2 for χ ,

$$\max(\text{dom } \phi) \wedge \chi(C) = \max(\text{dom } \phi) \wedge \chi(C'),$$

and similarly for the range of ϕ .

Generic refinability. $\phi \in \text{ParAut}(B)$ is *generically refinable within a class* $W \subseteq \text{ParAut}(B)$ if:

$$\begin{aligned}
& (\forall C, D \in \text{ACSAs}(B)) (\forall S \subseteq B^+, S \in L, \text{ with } S \text{ dense in } B^+) \\
& (\chi(C) \text{ compatible with } \max(\text{dom } \phi) \Rightarrow (\exists \phi' \in W, s \in S) \\
& (\phi' \text{ refines } \phi, \text{ and} \\
& \max(\text{ran } \phi') \leq s, \text{ and} \\
& \max(\text{dom } \phi') \wedge [C] \subseteq \text{dom } \phi', \text{ and} \\
& (\max(\text{ran } \phi') \leq \neg\chi(D), \text{ or } \max(\text{ran } \phi') \wedge [D] \subseteq \text{ran } \phi')).
\end{aligned}$$

Similarly, a filter $F \in \bigcup \Theta$ is *generically refinable in* (W, Θ) , where $W \subseteq \bigcup \Theta$ if:

$$\begin{aligned}
& (\forall D \in \text{ACSAs}(B)) (\forall M \in \Theta, S \subseteq B^+, S \in L, \text{ with } S \text{ dense in } B^+) \\
& (\exists N \in \Theta, F' \in N \cap W, s \in S \cap F') \\
& (M \in L(N) \text{ and } F' \text{ refines } F \text{ and} \\
& (\neg\chi(D) \in F', \text{ or } F' \cap D \text{ is an } L\text{-generic filter on } D)).
\end{aligned}$$

The set $\Gamma(\Theta)$ of fully refinable filters; the set Φ of fully refinable partial automorphisms

In the context of a particular χ , we use the above definitions to define a $\Gamma_\alpha(\Theta)$ hierarchy somewhat along the lines of $L_\alpha(\Theta)$, except here the sequences of levels shrink rather than grow:

$$\begin{aligned}
\Gamma_0(\Theta) &\equiv \{F \in \bigcup \Theta : F \text{ accords with } \chi\}; \\
\Gamma_{\alpha+1}(\Theta) &\equiv \{F \in \Gamma_\alpha : F \text{ is generically refinable in } (\Gamma_\alpha(\Theta), \Theta)\}; \\
\Gamma_\alpha(\Theta) &\equiv \bigcap_{\beta < \alpha} \Gamma_\beta(\Theta) \text{ for limit } \alpha; \\
\Gamma(\Theta) &\equiv \bigcap_{\alpha \in \text{Ord}} \Gamma_\alpha(\Theta).
\end{aligned}$$

We call $\Gamma(\Theta)$ the set of *fully generically-refinable* filters (in the context of a given χ and Θ); similarly, we define the set Φ of fully generically-refinable partial automorphisms:

$$\begin{aligned}
\Phi_0 &\equiv \{\phi \in \text{ParAut}(B) : \phi \text{ accords with } \chi\}; \\
\Phi_{\alpha+1} &\equiv \{\phi \in \Phi_\alpha : \phi \text{ is generically refinable within } \Phi_\alpha\}; \\
\Phi_\alpha &\equiv \bigcap_{\beta < \alpha} \Phi_\beta \text{ for limit } \alpha; \\
\Phi &\equiv \bigcap_{\alpha \in \text{Ord}} \Phi_\alpha.
\end{aligned}$$

Lemma 7.5 *Each $\Gamma(\Theta)$ -member is generically refinable in $(\Gamma(\Theta), \Theta)$; and each Φ -member is generically refinable within Φ .*

The $\Gamma_\alpha(\Theta)$ hierarchy must stop shrinking at some ordinal α at which $\Gamma(\Theta) = \Gamma_\alpha(\Theta) = \Gamma_{\alpha+1}(\Theta)$. For this α we have by definition

$$\Gamma_{\alpha+1}(\Theta) = \{F \in \Gamma_\alpha : F \text{ is generically refinable in } (\Gamma_\alpha(\Theta), \Theta)\};$$

replacing $\Gamma_\alpha(\Theta)$ and $\Gamma_{\alpha+1}(\Theta)$ with $\Gamma(\Theta)$ here yields the claim about $\Gamma(\Theta)$. The claim about Φ is proved the same way. \square

Lemma 7.6 *For all α , $\Phi_\alpha \in L$ and $\Gamma_\alpha(\Theta) \in L(\Theta)$; moreover, $\Phi \in L$ and $\Gamma(\Theta) \in L(\Theta)$.*

The proof is an application of our template for showing absoluteness of hierarchies, Lemmas 11.2 and 11.4. We show how to apply it to the $\Gamma(\Theta)$ hierarchy; the proof for Φ is similar. The template takes two predicates, $\Omega(\dots)$ and $\Psi(\dots)$, as its “inputs”. For the first, let $\Omega(x', \Theta)$ be “ $x' \in \bigcup \Theta$ and x' accords with χ ”; and for the second, let $\Psi(x'', (Y, \Theta, \beta))$ be “ $x'' \in Y \cap \bigcup \Theta$ and x'' is generically-refinable in $((Y \cap \bigcup \Theta), \Theta)$.”

We affirm first that our $\Omega(x', \Theta)$ is a Θ -absolute predicate, since the union operation is absolute and χ is a member of L . We affirm secondly that our $\Psi(x'', (Y, \Theta, \beta))$ is (Y, Θ, β) -absolute: this is because the definition of generic refinability for filters quantifies only over Θ , Y , and constructible subsets of B^+ .

With the claim about the $\Gamma_\alpha(\Theta)$ proved, the claim about $\Gamma(\Theta)$ follows from the observation that the levels $\Gamma_\alpha(\Theta)$ must stop shrinking when α reaches some ordinal δ , so that $\Gamma(\Theta) = \Gamma_\delta(\Theta)$. Similarly for Φ . \square

The next lemma establishes that $\Gamma(\Theta)$ is not empty, and in particular has at least one member of form $G \cap C$ from each Θ -member.

Lemma 7.7 *$(G \cap C) \in M \in \Theta \Rightarrow (G \cap C^{+\chi(C)}) \in M \cap \Gamma(\Theta)$.*

Suppose $(G \cap C) \in M \in \Theta$. Note that since $G \cap C \in \bigcup \Theta$, $\chi(C) \in G$, by definition of χ .

If ϕ denotes the identity function on C , then the one-element range refinement $\phi_{+\chi(C)}$ (defined above) copies $G \cap C$ to $G \cap C^{+\chi(C)}$; therefore $G \cap C^{+\chi(C)} \in M$ by Lemma 4.3.

$\chi(C^{+\chi(C)}) = \chi(C)$ by Lemma 7.4, so $\chi(C^{+\chi(C)}) \in G \cap C^{+\chi(C)}$; thus $(G \cap C^{+\chi(C)}) \in \Gamma_0(\Theta)$.

Note this holds for all C, M as in the lemma’s statement.

Now suppose there is a least ordinal $\delta > 0$ such that *some* C with $G \cap C \in \bigcup \Theta$ satisfies $(G \cap C^{+\chi(C)}) \notin \Gamma_\delta(\Theta)$, and fix one C witnessing this. Note δ must be a successor ordinal since otherwise $\Gamma_\delta(\Theta) = \bigcap_{\beta < \delta} \Gamma_\beta(\Theta)$, but $(G \cap C^{+\chi(C)}) \in \Gamma_\beta(\Theta)$ for all $\beta < \delta$.

Suppose that D, S, M witness $(G \cap C^{+\chi(C)})$ ’s violation of the definition of generic refinability in $(\Gamma_{\delta-1}(\Theta), \Theta)$. Fix any $s \in S \cap G$ such that $s \leq \chi(D)$, if $\chi(D) \in G$, or otherwise such that $s \leq \neg\chi(D)$. (Such an s exists because G is generic.) We now need to find $N \in \Theta$ and $F' \in N$ that together with s will satisfy the condition of definition of generic refinability for D, S, M , contradicting the supposed violation of that definition.

Let E be such that $G \cap E \in M$ (some such E exists by Lemma 4.3). $G \cap E \in \bigcup \Theta$ implies by definition of χ that $\chi(E) \in G$. Let E' denote $E^{+\chi(E)+s}$. By Lemma 7.4, $\chi(E') = \chi(E)$, so $\chi(E') \in G$.

We consider two cases.

If $\neg\chi(D) \in G$, let E'' denote $\overline{C^{+\chi(C)} \cup E'}$; then clause (ii) of Lemma 4.2 on Θ , plus the fact that the meet of any pair of G -members is a G -member, entails $\chi(E'') \in G$. So by definition of χ , $G \cap E'' \in \bigcup \Theta$. Let $F'' = G \cap E''$ and let N be the unique Θ -member such that $F'' \in N$. By the leastness of δ , $F'' \in \Gamma_{\delta-1}(\Theta)$. So our F'', N satisfy the generic-refinability requirement for $(G \cap C^{+\chi(C)}) \in \Gamma_\delta(\Theta)$, contradiction.

If $\chi(D) \in G$, we argue similarly, but with E'' defined as $\overline{C^{+\chi(C)} \cup D \cup E'}$. We know that the χ -values of the three subalgebras used to define E'' are all in G ; so as before we get $\chi(E'') \in G$, entailing $G \cap E'' \in \bigcup \Theta$. Letting $F'' = G \cap E''$ again, the same conclusions about F'' follow, and in addition we have that $F'' \cap D$ is a generic filter on D , fulfilling the last clause of generic refinability. So again F'', N satisfy the generic-refinability requirement for $(G \cap C^{+\chi(C)}) \in \Gamma_\delta(\Theta)$.

Thus no such ordinal δ can exist, and $(G \cap C^{+\chi(C)}) \in \Gamma(\Theta)$. \square

7.3 The self-collection condition in terms of filters

Theorem 7.8 (Self-collection condition in terms of filters) *Under Stipulation 3.11, $G \in L(\Theta)$ iff there exists $D \in \text{ACSAs}(B)$ such that $(G \cap D) \in \Gamma(\Theta)$ and all $(G \cap D)$'s refinements in $\Gamma(\Theta)$ are mutually compatible, in which case the proposition “ $x \in F \in \Gamma(\Theta)$ for some refinement F of $G \cap D$ ” serves as the predicate $\Psi(\dots)$ for Lemma 5.1.*

For $D \in \text{ACSAs}(B)$ satisfying $(G \cap D) \in \Gamma(\Theta)$, let Q_D be the set of all fully-refinable refinements of $G \cap D$, that is:

$$Q_D \equiv \{F \in \Gamma(\Theta) : G \cap D \subseteq F\}.$$

Since $\Gamma(\Theta) \in L(\Theta)$ (Lemma 7.6), $Q_D \in L(\Theta)$, and $\bigcup Q_D \in L(\Theta)$.

We begin with the “if” direction. If D exists as in the lemma’s statement, then all Q_D -members are mutually compatible; we will show this entails $G = \bigcup Q_D$, which suffices. Note first that by Lemma 7.7, $G \cap D^{+\chi(D)}$ is a refinement of $G \cap D$ that is a member of $\Gamma(\Theta)$. Now for any $b \in B$, consider $(D^{+\chi(D)})^{+b}$. By Lemma 7.4, we have

$$\chi((D^{+\chi(D)})^{+b}) = \chi(D^{+\chi(D)}) = \chi(D);$$

thus $(D^{+\chi(D)})^{+b}$ already has its own χ -value as a member, and so again by Lemma 7.7, $(G \cap (D^{+\chi(D)})^{+b})$ is a refinement of $G \cap D$ in $\Gamma(\Theta)$, and so is a Q_D -member. So $b \in G$ iff $b \in \bigcup Q_D$, and since this holds for all b , $G = \bigcup Q_D$.

Next we prove the “only if” direction; supposing there is no D meeting the condition stated in the lemma, we will show that Θ does not construct G . Suppose towards a contradiction that Θ *did* construct G , and let Ψ , C , and \vec{c} witness this in Lemma 5.1. Some $g \in G$ must force them to be witnesses; so let us choose $g \in G, g \leq \chi(C)$ such that

$$g \Vdash \{x : \Psi(x, \dot{\Theta}, \dot{G} \cap \check{C}, \vec{c})\} = \dot{G}.$$

We will now show that our suppositions permit the existence of a G' contradicting Lemma 5.2.

Let D be $(C^{+g})^{+\chi(C)}$. Note that $G \cap D \in \Gamma(\Theta)$ by Lemma 7.7. We will obtain our G' by forcing over the set Q_D ordered by reverse inclusion. Note that $L(G)$ is the ground model for this forcing. Let Y be a generic filter obtained by this forcing, and define G' to be $\bigcup Y$.

G' is clearly a filter on B^+ . It is generic over L by the following argument. Let $S \in L$ be any dense subset of B^+ . Because each $F \in Q_D$ satisfies $F \in \Gamma(\Theta)$, and each of its refinements in $\Gamma(\Theta)$ is in Q_D , any $F \in Y$ can be refined (by Lemma 7.5) to $F' \in Q_D$ with $F' \cap S \neq \emptyset$. Thus, by genericity, some such F' is in Y , and $G' \cap S \neq \emptyset$.

It follows immediately from the definitions of D and of Q_D that $G \cap D = G' \cap D$; and since $g \in G \cap D$, also $g \in G' \cap D$.

To show that $G' \neq G$, note every $F \in Q_D$ can be refined to some $F' \in Q_D$ that is incompatible with G . For if some F were not so refinable, it would equal $G \cap D'$ for some $D' \supseteq D$, and all its refinements in $\Gamma(\Theta)$ would be mutually compatible (being compatible with G); D' would thus meet the requirements to be the D that we are supposing not to exist. By genericity/density, some F' inconsistent with G must therefore be in Y , whence $G' \neq G$.

To see that G' gives the same interpretation as G to $\dot{\Theta}$, consider first any unconstructible $M \in (\dot{\Theta})_G$. (For the remainder of this argument we use the notation $(\dot{\Theta})_G$ for G 's interpretation of $\dot{\Theta}$, which we had been calling Θ , in order to avoid confusing it with $(\dot{\Theta})_{G'}$.) We will show that also $M \in (\dot{\Theta})_{G'}$.

Let $F \in Q_D$ be arbitrary. Since $F \in \Gamma(\Theta)$, there exists by Lemma 7.5 and the definition of generic refinability some $N \in (\dot{\Theta})_G$, and $F' \in N \cap \Gamma(\Theta)$, such that F' refines F and $M \in L(N)$. Since any $F \in Q_D$ is refinable to such an F' , some such F' must, by Y 's genericity, be a member of Y . Let $E \in \text{ACSA}_s(B)$ be the subalgebra on which F' is an ultrafilter. The definition of χ -accordance of filters along with $G' = \bigcup Y$ gives us $\chi(E) \in G'$; thus F' is a member of $\bigcup (\dot{\Theta})_{G'}$. Now since $M \in L(N)$, it follows from Lemma 4.7, the definition of Θ , and Lemma 4.3 that there must be an ACSA $E' \subseteq E$ such that $G' \cap E' \in M$. Lemma 6.2 applied to $(\dot{\Theta})_{G'}$ then entails that $G' \cap E'$ is a member of $\bigcup (\dot{\Theta})_{G'}$, so $M \in (\dot{\Theta})_{G'}$.

Conversely, fix an unconstructible $M \in (\dot{\Theta})_{G'}$. Since G' , like G , is an L -generic ultrafilter on B , Lemma 7.7 applies to it, ensuring the existence of an E such that $G' \cap E \in M$. Now it would suffice to show that there exists an $F' \in Y$ such that F' refines $G' \cap E$, because then:

$$F' \in Y \Rightarrow$$

$F' \in Q_D \Rightarrow$
 $F' \in \bigcup (\dot{\Theta})_G \Rightarrow$
 $F' \in N \in (\dot{\Theta})_G$ for some $N \Rightarrow$
 F' is a copy of some $G \cap A \in N$ via some $\phi \in \text{ParAut}(B)$ (Lemma 4.3) \Rightarrow
 $G' \cap E$ is a copy via (a restriction of) ϕ of $G \cap A'$ for some $A' \subseteq A \Rightarrow$
 $G' \cap E \in \bigcup (\dot{\Theta})_G$ (Lemma 6.2) \Rightarrow
 $M \in (\dot{\Theta})_G$.

To see that such an $F' \in Y$ does exist, consider that by the definition of generic refinability, any $F \in Q_D$ can be refined to some $F' \in Q_D$ such that either $\neg\chi(E) \in F'$, or $F' \cap E$ is a generic filter on E . By Y 's genericity, then, some such F' is in Y , and since we already know $\chi(E) \in \bigcup Y$, the second alternative must hold, which entails that F' refines $G' \cap E$. \square

8 Self-Collection Condition In Terms Of Isomorphisms

The preceding theorem gives a self-collection condition that can be checked in the generic extension, stated in terms of generic filters; we now prove lemmas that will let us translate this condition into a condition on the boolean algebra B (given in Theorem 8.7) that can be checked in the ground model, stated in terms of $\text{ParAut}(B)$.

Definition. $\phi \in \text{ParAut}(B)$ fixes $C \in \text{ACSAs}(B)$ if it is a refinement of the identity function on C . This definition is equivalent to:

$$\max(\text{dom } \phi) \wedge [C] \subseteq \text{dom } \phi, \text{ and } (\forall b \in \text{dom } \phi)(\phi(b) \uparrow C = b \uparrow C).$$

Lemma 8.1 *If ϕ copies $G \cap C$ to F and $b \in G$, ϕ^{+b} copies $G \cap C^{+b}$ to F .*

(We assume ϕ^{+b} is well defined, so $b \leq \max(\text{dom } \phi)$.) Let D denote the ACSA on which F is an ultrafilter. We must show that for all $d \in D$,

$$(*) \quad (\exists c \in G \cap \text{dom } \phi)(\phi(c) \leq d) \iff (\exists c' \in G \cap \text{dom } \phi^{+b})(\phi^{+b}(c') \leq d).$$

We showed in Lemmas 7.2 and 7.3 that the map $\psi : c \mapsto b \wedge c$ defined on $C \upharpoonright (b \uparrow C)$ is an isomorphism, and that $\phi^{+b}(c') = \phi(\psi^{-1}(c'))$ for all $c' \in \text{dom } \phi^{+b}$. Since G is an ultrafilter on B and $b \in G$, we have for all $c \in \text{dom } \phi$ that $c \in G \iff c \wedge b \in G$. Now every $c' \in \text{dom } \phi^{+b}$ is of form $c \wedge b$ for some $c \in \text{dom } \phi$, from which (*) follows. \square

Lemma 8.2 *If ϕ copies $G \cap C$ to F , and F' refines F , and $F' \in \theta(Y)$ for some singly-generated continuum $Y \in L[G]$, there exist ϕ' and $D' \supseteq C$ such that ϕ' copies $G \cap D'$ to F' and ϕ' refines ϕ .*

By Lemma 4.3, *some* $\gamma \in \text{ParAut}(B)$ copies *some* $G \cap C'$ to F' . Let $E \subseteq E'$ be the ACSAs on which F, F' are generic ultrafilters respectively. Let $D \subseteq C'$ be an ACSA of which $\gamma^{-1}[E \cap \text{ran } \gamma]$ is a principal ideal, so that $D \cap G$ is interconstructible with $F' \cap E = F$. Thus $D =_G C$.

By Lemma 4.7 applied “in both directions,” there exist $g_1, g_2 \in G$ such that $g_1 \wedge [D] \subseteq g_1 \wedge [C]$ and $g_2 \wedge [C] \subseteq g_2 \wedge [D]$. Since G is a filter, $g \equiv g_1 \wedge g_2 \in G$, and $g \wedge [C] = g \wedge [D]$.

Define $\gamma' \equiv \gamma^{+g}$, the one-element refinement of γ by g . If we define $D' \equiv \overline{C \cup C'}$, it is easy to check that $\text{dom } \gamma'$ is a principal ideal of D' , and that γ' refines ϕ , and copies $G \cap D'$ to F' . \square

Corollary 8.3 *If $F' \in \theta(Y)$ for some singly-generated continuum Y , and $G \cap C \subseteq F'$ for some $C \in \text{ACSAs}(B)$, then there exists $D' \supseteq C$ and $\phi' \in \text{ParAut}(B)$ such that ϕ' copies $G \cap D'$ to F' , and ϕ' fixes C .*

In Lemma 8.2, let F be $G \cap C$, and let ϕ be the identity function on C . \square

Lemma 8.4 *In light of Stipulation 3.11’s consequences for χ (Lemma 6.1), every $F \in \Gamma(\Theta)$ is the copy of some $G \cap E$ via some $\phi \in \Phi$.*

In $L[G]$, by Lemma 4.3, every generic filter F on any $D \in \text{ACSAs}(B)$ is a copy via some $\phi \in \text{ParAut}(B)$ of $G \cap E$ for some E . We must show that when $F \in \Gamma(\Theta)$, the ϕ and E can be chosen so that $\phi \in \Phi$. In fact we will show something stronger: given any choice of ϕ, E , there exists b such that $\phi^{+b} \in \Phi$ and ϕ^{+b} copies $G \cap E^{+b}$ to F .

Let $\Delta(\alpha)$ be the following assertion about an ordinal α :

$\Delta(\alpha) \iff$ *For every $F \in \Gamma(\Theta)$ and every ϕ copying some $G \cap E$ to F , there exists $b \in G$, $b \leq \max(\text{dom } \phi)$, such that $\phi^{+b} \in \Phi_\alpha$.*

Any such ϕ^{+b} will copy $G \cap E^{+b}$ to F , by Lemma 8.1. Thus our “stronger” claim above will follow if $\Delta(\alpha)$ holds for all α (since $\Phi = \Phi_\alpha$ for some α). Let us suppose towards a contradiction that $\Delta(\alpha)$ fails for some least α . Fix F, ϕ , and E witnessing the failure, and let D be the ACSA on which F is an ultrafilter. There are then three cases.

Case I: $\alpha = 0$. Since $F \in \Gamma_0(\Theta)$, it accords with χ , so $\chi(D)$ is in F ’s upwards closure in B . Choose some $e \in \phi[G \cap \text{dom } \phi]$ with $e \leq \chi(D)$. Let $b = \phi^{-1}(e)$. Then $\phi \upharpoonright b = \phi^{+b}$, and $b \in G$. Now since $F \in \bigcup \Theta$, $G \cap E \in \bigcup \Theta$ and so $\chi(E) \in G$; thus $g \equiv \chi(E) \wedge b$ is in G , since G is a filter. It is then easy to check that ϕ^{+g} accords with χ , so $\phi^{+g} \in \Phi_0$. This contradicts the supposition that F, ϕ, E witness $\neg \Delta(0)$.

Case II: α is a successor ordinal. Since F, ϕ , and E supposedly witness $\neg \Delta(\alpha)$, we have for all $b \in G$ with $b \leq \max(\text{dom } \phi)$ that $\phi^{+b} \notin \Phi_\alpha$. By the leastness of α , however, there does exist such a b satisfying $\phi^{+b} \in \Phi_{\alpha-1}$. Indeed, whenever $b \in G$ and $b \leq \max(\text{dom } \phi)$, ϕ^{+b} will copy $G \cap E^{+b}$ to F (by Lemma 8.1), and $\Delta(\alpha-1)$ will hold for F, ϕ^{+b} , and E^{+b} too,

so that there will exist $b' \leq b$ satisfying $\phi^{+b'} \in \Phi_{\alpha-1}$. For all such b' , $\phi^{+b'}$ is in $\Phi_{\alpha-1}$ but fails to be generically refinable within $\Phi_{\alpha-1}$, and this failure must be witnessed by some C, D, S as in the definition of generic refinability (for isomorphisms); in particular $b' \wedge \chi(C) \neq 0$. Now by genericity we must have $b' \wedge \chi(C) \in G$ for some such b' having associated witnesses C, D, S (note this implies $\chi(C) \in G$). Fix such a b' and associated C, D, S .

Now define $C' \equiv \overline{C \cup E}$. Because $\chi(C), \chi(E) \in G$, stipulation (ii) on χ (from Lemma 6.1) entails $\chi(C') \in G$. Therefore there exists $M \in \Theta$ such that $G \cap C' \in M$. Since $F \in \Gamma(\Theta)$, Lemma 7.5 ensures that it refines generically in $(\Gamma(\Theta), \Theta)$. In particular (reading off the definition of generic refinability for filters) F refines to some $F' \in N \in \Theta$, for some N with $M \in L(N)$, such that $F' \cap S \neq \emptyset$, and $F' \in \Gamma(\Theta)$, and either $\neg\chi(D) \in F'$ or $F' \cap D$ is an L -generic filter on D .

Now by Lemma 8.2, this $F' \in \Gamma(\Theta)$ is a copy of $G \cap C''$ for some $C'' \supseteq C'$, via some ϕ' that refines ϕ . We know $\Delta(\alpha - 1)$ holds, so some ϕ'^{+e} with $e \in G$ and $e \leq \max(\text{dom } \phi')$ satisfies $\phi'^{+e} \in \Phi_{\alpha-1}$. By Lemma 8.1, ϕ'^{+e} copies $C''^{+e} \cap G$ to F' . Consider the restriction ϕ'' of ϕ'^{+e} to some principal ideal of C''^{+e} such that $\max(\text{ran } \phi'')$ is \leq some member of $F' \cap S$. It is then straightforward to check that ϕ'' is a refinement of $\phi^{+b'}$ that satisfies the requirements of generic refinability within $\Phi_{\alpha-1}$ for our fixed C, D and S , contradicting our choices.

Case III: α is a limit ordinal. The argument here is similar. Recall that for limit α , Φ_α is just the intersection of the Φ_β 's with $\beta < \alpha$. Recall our fixed F, ϕ , and E witnessing $\neg\Delta(\alpha)$: again, for all $b \in G$ with $b \leq \max(\text{dom } \phi)$, $\phi^{+b} \notin \Phi_\alpha$. But now since α is a limit ordinal, this means for all such b there exists $\gamma < \alpha$, and thus a *least* $\gamma < \alpha$, such that $\phi^{+b} \notin \Phi_\gamma$; moreover this γ must be a successor ordinal. As before, for each such b , some C, D, S will witness $\phi^{+b} \in \Phi_{\gamma-1} \setminus \Phi_\gamma$ (for b 's particular γ), so that in particular $b \wedge \chi(C) \neq 0$. As before, we have by genericity that $b \wedge \chi(C) \in G$ for some such b with its associated C, D, S . The contradiction is then obtained just as before, but with γ in place of α . \square

Lemma 8.5 *If the F in Lemma 8.4 satisfies $G \cap D \subseteq F$ for some D , then the E and ϕ can be chosen such that ϕ fixes D .*

The proof of Lemma 8.4 showed that, given any choice of ϕ, E such that ϕ copies $G \cap E$ to F , there exists b such that $\phi^{+b} \in \Phi$ and ϕ^{+b} copies $G \cap E^{+b}$ to F . By Corollary 8.3, we may choose a ϕ that fixes D (is a refinement of the identity function on D), and the resulting ϕ^{+b} will also be a refinement of the identity on D . \square

Lemma 8.6 *If $\phi \in \Phi$, $\max(\text{dom } \phi) \in G$, and ϕ copies $(G \cap \text{dom } \phi)$ to F , then $F \in \Gamma(\Theta)$.*

Note this is a converse to Lemma 8.4; in this direction, we will argue much the same way. Let $\Psi(\alpha)$ be the following assertion about an ordinal α :

$\Psi(\alpha) \iff$ every filter F that is a copy of $G \cap E$ for some E , via some $\phi \in \Phi$, is a member of $\Gamma_\alpha(\Theta)$.

Suppose toward a contradiction that $\Psi(\alpha)$ fails for some least α . Fix F, E, ϕ witnessing this failure, and let D denote the ACSA on which F is an ultrafilter.

Case I: $\alpha = 0$. Since ϕ accords with χ , $\max(\text{ran } \phi) \leq \chi(D)$. Since $\max(\text{dom } \phi) \in G$, $\max(\text{ran } \phi) \in \phi[G \cap \text{dom } \phi] \subseteq F$, so $\chi(D)$ is in the upward closure of F , which is just the definition of accordance with χ for F . Also $F \in \bigcup \Theta$ follows from Lemma 4.3 because $G \cap E \in \bigcup \Theta$ (since $\max(\text{dom } \phi) \in G$ and $\max(\text{dom } \phi) \leq \chi(E)$). So $F \in \Gamma_0(\Theta)$ after all.

Case II: α is a successor ordinal. By α 's leastness, $F \in \Gamma_{\alpha-1}(\Theta)$; but we have supposed $F \notin \Gamma_\alpha(\Theta)$, so F fails to refine generically in $(\Gamma_{\alpha-1}(\Theta), \Theta)$. Let some D, S, M witness this failure, as in the definition of generic refinability of filters. In particular note that $M \in \Theta$; since also $G \cap E \in \bigcup \Theta$, there must exist by the directedness of $\bigcup \Theta$ (stipulation (ii) on Θ from Lemma 4.2) some $N \in \Theta$ and $C \in \text{ACSAs}(B)$ such that $E \subseteq C$, and $G \cap C \in N$, and $M \in L(N)$. Fix such an N and C . Since $G \cap C \in \bigcup \Theta$, $\chi(C) \in G$, so $\chi(C)$ is compatible with $\max(\text{dom } \phi)$ (which is also in G). Now apply the definition of ϕ 's being generically refinable within Φ , using our present C, D and S , to obtain $s \in S$ and $\phi' \in \Phi$ refining ϕ , meeting the several requirements of that definition.

Let $A \in \text{ACSAs}(B)$ have $\text{ran } \phi'$ as a principal ideal. Then A^{+s} also has it as a principal ideal, because $\max(\text{ran } \phi') \leq s$. Now consider the two cases in the final clause of the definition that was just invoked to obtain ϕ' . If $\max(\text{ran } \phi) \leq \neg\chi(D)$, define

$$A' \equiv (A^{+s})^{+\neg\chi(D)};$$

otherwise we define

$$A' \equiv \overline{A^{+s} \cup D}.$$

In both cases we still have $\text{ran } \phi$ a principal ideal of A'' ; in the latter case this is because $\max(\text{ran } \phi') \wedge [D] \subseteq \text{ran } \phi'$.

Now note that $F' \in \Gamma_{\alpha-1}(\Theta)$, by the way F' was obtained as a copy via $\phi' \in \Phi$ and because $\Psi(\alpha - 1)$ holds. It is then straightforward to verify, by $F' \in \Gamma_{\alpha-1}(\Theta)$, by the requirements that ϕ' meets, and by the definition of A'' , that our N, F' , and s meet all the requirements of the generic refinability definition for our D, S, M — the requirements that we supposed unsatisfiable.

Case III: There is no Case III here because the least α at which $\Psi(\alpha)$ fails cannot be a limit ordinal: if every filter F of the kind specified in the definition of $\Psi(\alpha)$ is a member of $\Gamma_\beta(\Theta)$ for all $\beta < \alpha$, then $F \in \Gamma_\alpha(\Theta)$, since the latter is just the intersection of all the $\Gamma_\beta(\Theta)$ with $\beta < \alpha$. \square

Theorem 8.7 *Let X be the set of all $d \in B$ for which there exists $D \in \text{ACSAs}(B)$, $d \leq \chi(D)$, such that every $\phi \in \Phi$ that fixes D and satisfies $\max(\text{dom } \phi) \wedge d > 0$ is an identity mapping. Then under Stipulation 3.11, $\|G \in L(\Theta)\| = \bigvee X$.*

It suffices to show that there exists a $d \in X \cap G$ just if a D meeting the condition of Theorem 7.8 exists.

Suppose D witnesses $d \in X$, for some $d \in G$. Thus we have $d \leq \chi(D)$, so that $\chi(D) \in G$ and $D \cap G \in \bigcup \Theta$, and every $\phi \in \Phi$ that fixes D and satisfies $\max(\text{dom } \phi) \wedge d > 0$ is an identity mapping. We claim that $D^{+\chi(D)}$ satisfies the condition of Theorem 7.8. We must show (reading off that condition) that $G \cap D^{+\chi(D)} \in \Gamma(\Theta)$ and all refinements $F \in \Gamma(\Theta)$ of $G \cap D^{+\chi(D)}$ are mutually compatible. The first requirement holds by Lemma 7.7. For the second, suppose that some $F, F' \in \Gamma(\Theta)$ violated it and (without loss of generality) that F is not compatible with G . By Lemma 8.4, F would have to be a copy, via some non-identity $\phi \in \Phi$, of $G \cap E$, for some E . By Lemma 8.5, we can also require that ϕ fix $D^{+\chi(D)}$, and hence D . Since ϕ copies $G \cap E$, we have $\max(\text{dom } \phi) \in G$; thus $\max(\text{dom } \phi) \wedge d$ belongs to G and so must be > 0 . This ϕ contradicts our assumption that D witnessed $d \in X$.

Similarly, in the other direction, suppose D satisfying $(G \cap D) \in \Gamma(\Theta)$ meets the condition of Theorem 7.8: every $F \in \Gamma(\Theta)$ that refines $G \cap D$ is of form $G \cap E$ for some $E \supseteq D$. Choose some $d \in G$ that forces this assertion to hold. We claim that D witnesses $d \in X$. If not, some non-identity $\phi \in \Phi$ fixes D and satisfies $\max(\text{dom } \phi) \wedge d > 0$. But by Lemma 8.6, if we had $\max(\text{dom } \phi) \in G$, ϕ would copy $(G \cap \text{dom } \phi)$ to a filter $F \in \Gamma(\Theta)$. Since ϕ is not an identity mapping, this F is not of form $G \cap E$. So $\max(\text{dom } \phi) \wedge d$ would force this F to violate the assertion that d supposedly forced. \square

9 Negative Results On Self-Construction

We will now establish that the self-collection criteria derived in the preceding sections are not trivial to satisfy. In particular, we will show that any \mathcal{N} meeting Stipulation 3.11 fails to self-collect into $\mathbb{R}(G)$ if B has a “flexible homogeneity” property ensuring that it has certain partial automorphisms. Since this property makes no mention of \mathcal{N} (nor of any structures defined from it, like Θ or χ), one upshot is that, even assuming B yields a rich structure of continua when used for forcing, it may just have too many partial automorphisms, so that no amount of cleverness in designing a B -name for \mathcal{N} can suffice to make \mathcal{N} self-collect into $\mathbb{R}(G)$. We will show that this is the case when B is a Cohen-real or random-real algebra.

9.1 A homogeneity property that precludes self-collection

We now begin arguments culminating with Theorem 9.6, to show that when our boolean algebra B has the “flexible homogeneity” property defined just below, B cannot yield singly-generated continua that self-collect into another such continuum.

Definitions:

The definitions and stipulations of Sections 2, 3, and 4 remain in force (for B , G , ACSAs(B), \mathcal{N} , Θ , χ , etc.).

B is *homogeneous* if it is isomorphic to each of its principal ideals.

B is *ACSA-homogeneous* if it is isomorphic to every principal ideal of every $C \in \text{ACSAs}(B)$.

B is *flexibly homogeneous* if:

- (1) B is ACSA-homogeneous;
- (2) there exists $C \in \text{ACSAs}(B)$ satisfying $1 \Vdash C \not\leq_G B$;
- (3) for each C satisfying (2), $C \subseteq B$ is not a rigid boolean inclusion (see Section 6);
- (4) each isomorphism between any C, C' satisfying (2) extends to an automorphism of B .

We will also apply the above terms to a principal ideal $B \upharpoonright b$ if it meets the relevant definitions when considered as a boolean algebra in its own right (with greatest element b and negation operation derived from B 's negation as $x \mapsto b \wedge \neg x$).

Lemma 9.1 *If B is flexibly homogeneous, then each principal ideal of each $C \in \text{ACSAs}(B)$ (including the “improper ideal,” C itself) is also flexibly homogeneous.*

Immediate from clause (1) of flexible homogeneity. \square

Lemma 9.2 *Given $C \in \text{ACSAs}(B)$ and $b \in B^+$ such that $b \leq \chi(C)$, and assuming that $\|\mathcal{N} \text{ self-collects into } \mathbb{R}(G)\| = 1$, there exists $E \supset C$ such that*

$$b \wedge \chi(E) \wedge \|E >_G C\| > 0,$$

and each term in that conjunction is a member of E .

The main point here is that no continuum of form $\mathbb{R}(G \cap C)$ can be maximal among continua of that form contained in $\bigcup \mathcal{N}$; there must exist a strictly larger $\mathbb{R}(G \cap E) \subseteq \bigcup \mathcal{N}$.

The key is to show that some $D \in \text{ACSAs}(B)$ must satisfy

$$b \wedge \chi(D) \wedge \|D \not\leq_G C\| > 0.$$

Suppose not. Then for all D ,

$$(b \wedge \chi(D)) \Vdash D \leq_G C.$$

From this and the definitions of χ and Θ it follows that

$$b \Vdash (\mathbb{R}(G \cap D) \subseteq \bigcup \mathcal{N}) \Rightarrow (\mathbb{R}(G \cap D) \subseteq \mathbb{R}(G \cap C)),$$

and since that holds for all D , we have

$$b \Vdash \mathbb{R}(\bigcup \mathcal{N}) = \mathbb{R}(G \cap C).$$

From this and $\|\mathcal{N}$ self-collects into $\mathbb{R}(G)\| = 1$ it follows that

$$b \Vdash \mathbb{R}(G \cap C) = \mathbb{R}(G).$$

Now since $b \leq \chi(C)$, it follows from the definitions of Θ and χ in terms of \mathcal{N} that

$$b \Vdash (\exists C')(\mathbb{R}(G \cap C) \subseteq \mathbb{R}(G \cap C') \text{ and } \mathbb{R}(G \cap C') \in \mathcal{N}),$$

which along with the previous step gives us

$$b \Vdash \mathbb{R}(G) \in \mathcal{N},$$

which would violate clause (i) of the definition of self-collection.

The foregoing permits us to fix an ACSA D such that when we define

$$e \equiv b \wedge \chi(D) \wedge \|D \not\leq_G C\|,$$

we have $e > 0$.

Now define $E = \overline{C \cup D}$. Plainly $C \subseteq E$, $C \leq_G E$, and $D \leq_G E$. Since $e \leq \chi(D)$, and $e \leq b \leq \chi(C)$, we have by (ii) of Lemma 6.1 that $e \leq \chi(E)$. We claim that $e \leq \|E >_G C\|$. This follows from the fact (Corollary 3.3) that $<_G$ is guaranteed with boolean value 1 to be a preorder: if some $e' \leq e$ forced $E =_G C$ and we had $e' \in G$, then in $L[G]$ we would have $E =_G C$ and $D \leq_G E$, entailing $D \leq_G C$, contradicting our choice of D .

Thus E meets all our requirements except possibly the requirement (needed for technical simplification later) that b , $\chi(E)$ and $\|E >_G C\|$ be members. To ensure this, simply replace E with $E^{+b+\chi(E)+\|E>_G C\|}$; Lemma 7.4 ensures that E 's χ -value stays the same, and clearly the other requirements on E in the statement of the lemma will still hold when E is enlarged. \square

Lemma 9.3 *Suppose that B is flexibly homogeneous and that $\|\mathcal{N}$ self-collects into $\mathbb{R}(\dot{G})\| = 1$; if $C \in \text{ACSAs}(B)$ and $g \in G$ were witnesses to \mathcal{N} 's self-collection into $\mathbb{R}(G)$ as in Theorem 8.7, then there would exist $\phi \in \Phi_0$ with $\max(\text{dom } \phi) \leq g$ that fixes C but is not an identity function.*

Note the fact that C, g witness Theorem 8.7 entails $g \leq \chi(C)$. Consider $B \upharpoonright g$ and $C^{+g} \upharpoonright g$. By Lemma 9.1 they are flexibly homogeneous. By Lemma 9.2, there exists $E \in \text{ACSAs}(B \upharpoonright g)$ such that $E \supset (C^{+g} \upharpoonright g)$, and when we define

$$e \equiv \chi(E) \wedge \|E >_G (C^{+g} \upharpoonright g)\|,$$

we have $0 < e \leq g$, and also $e \in E$.

By Lemma 9.1 $E \upharpoonright e$ is flexibly homogeneous; in particular it satisfies clause (3), so there exists a nontrivial automorphism ϕ of $E \upharpoonright e$ whose restriction to $C^{+e} \upharpoonright e$ is the identity. And we have $\phi \in \Phi_0$ because $\max(\text{ran } \phi) = \max(\text{dom } \phi) = e \leq \chi(E')$, where E' is any ACSA of (all of) B such that $E' \upharpoonright e = E \upharpoonright e$. \square

Lemma 9.4 *If $\Phi_1 = \Phi_0$ then $\Phi = \Phi_0$.*

Suppose $\Phi_1 = \Phi_0$ but there is a least ordinal α (necessarily > 1) such that there exists $\phi \in \Phi_0 \setminus \Phi_\alpha$. Note α cannot be a limit ordinal since in that case Φ_α would be the intersection of the Φ_β for all $\beta < \alpha$. Thus there exists some $\phi \in \Phi_{\alpha-1}$ that fails to refine generically within $\Phi_{\alpha-1}$. However $\Phi_{\alpha-1} = \Phi_0$ and we already know ϕ refines generically within Φ_0 , since $\phi \in \Phi_1$. \square

Lemma 9.5 *Suppose B is flexibly homogeneous; if $\|\dot{\mathcal{N}} \text{ self-collects into } \mathbb{R}(\dot{G})\|$ were equal to 1, then $\Phi = \Phi_0$.*

By Lemma 9.4 it suffices to show that any $\phi \in \Phi_0$ satisfies $\phi \in \Phi_1$ — equivalently, refines generically within Φ_0 . So fix $\phi \in \Phi_0$, and, for the parameters of the definition of generic refinability, fix an arbitrary dense subset $S \subseteq B^+$ and $C, D \in \text{ACSAs}(B)$, such that when we define

$$a \equiv \chi(C) \wedge \max(\text{dom } \phi),$$

we have $a > 0$. We will show that the ϕ' required in the definition does exist.

Let $C_0 \in \text{ACSAs}(B)$ be such that $\text{dom } \phi$ is a principal ideal of C_0 , and likewise for D_0 with $\text{ran } \phi$. (In general, variable names involving “C” will denote ACSA’s involved with the domain of the ϕ' we are seeking, and likewise with “D” and the range of ϕ' .) Note that since $\phi \in \Phi_0$, we have $\max(\text{dom } \phi) \leq \chi(C_0)$ and $\max(\text{ran } \phi) \leq \chi(D_0)$.

Define $C_1 \equiv \overline{C_0 \cup C}$. By clause (ii) of Lemma 6.1, $\chi(C_1) = \chi(C_0) \wedge \chi(C)$. Therefore we have $a \leq \chi(C_1)$.

By Lemma 9.2, there exists $E_C \supseteq C_1$ such that, when we set

$$c \equiv a \wedge \chi(E_C) \wedge \|E_C >_G C_1\|,$$

we have $c > 0$, and each conjunct in c ’s definition is a member of E_C .

Now ϕ^{+c} , the one-element domain refinement of ϕ by c , is well-defined because $c \leq a \leq \max(\text{dom } \phi)$. Note $\max(\text{dom } \phi^{+c}) = c$, and

$$\max(\text{ran } \phi^{+c}) \leq \max(\text{ran } \phi) \leq \chi(D_0).$$

To fix the ACSA E_D of which $\text{ran } \phi'$ will be a principal ideal, we consider two cases.

Case I: $(\chi(D) \wedge \max(\text{ran } \phi^{+c})) > 0$.

In this case let s be any S -member $\leq (\chi(D) \wedge \max(\text{ran } \phi^{+c}))$, and define

$$D_1 \equiv \overline{D_0 \cup D}.$$

By (ii) of Lemma 6.1, $\chi(D_1) = \chi(D_0) \wedge \chi(D)$. Also we have

$$s \leq \chi(D_1) = \chi(D_0) \wedge \chi(D),$$

because

$$s \leq \max(\text{ran } \phi^{+c}) \leq \chi(D_0).$$

Case II: $\chi(D) \wedge \max(\text{ran } \phi^{+c}) = 0$.

In this case let s be any S -member below $\max(\text{ran } \phi^{+c})$, and let $D_1 = D_0$ (so again we have $s \leq \chi(D_1)$).

Conclusion of argument for both cases:

Now use Lemma 9.2 to obtain $E_D \supseteq D_1$ such that when we define

$$d \equiv s \wedge \chi(E_D) \wedge \|E_D >_G D_1\|,$$

we have $d > 0$, and each conjunct in d 's definition is a member of E_D .

Consider the isomorphism $(\phi^{+c})_{+d}$. Let C'_0 denote its domain, and let D'_0 denote its range. Let c' denote $\max(\text{dom}((\phi^{+c})_{+d}))$; note $C'_0 = c' \wedge [C_0]$.

Let $E'_C \equiv E_C \upharpoonright c'$, and $E'_D \equiv E_D \upharpoonright d$. We consider E'_C and E'_D as boolean algebras in their own right. It is easily verified that $C'_0 \in \text{ACSAs}(E'_C)$ and $D'_0 \in \text{ACSAs}(E'_D)$.

Since $C_1 \supseteq C_0$, $\|C_1 \geq_G C_0\| = 1$, so $c' \leq \|E_C >_G C_0\|$. Thus the boolean 1 of E'_C forces $E'_C >_G C'_0$.

Now E'_C is isomorphic to E'_D by ACSA-homogeneity of B ; choose an isomorphism $\xi : E'_C \rightarrow E'_D$ witnessing this. Consider the map $\xi \circ ((\phi^{+c})_{+d})^{-1}$ with domain D'_0 . Its range is some ACSA E^* of E'_D . Since C'_0 is the range of $((\phi^{+c})_{+d})^{-1}$, and the boolean 1 of E'_C forces $E'_C >_G C'_0$, the boolean 1 of E'_D must (by the fact that ξ is an isomorphism) force $\|E^* <_G E'_D\|$.

Consider the inverse of $\xi \circ ((\phi^{+c})_{+d})^{-1}$, which maps E^* onto D'_0 . By clause (4) of flexible homogeneity, it extends to an automorphism ζ of E'_D . Now define $\phi' \equiv \zeta \circ \xi$ by composition. It is straightforward to check that ϕ' is the refinement of ϕ demanded by the definition of generic refinability for our chosen parameters S, C, D . \square

Theorem 9.6 *If $B \in L$ is a countably-completely-generated boolean algebra that is flexibly homogeneous, and G is a generic ultrafilter on B , and $x \in \mathbb{R}(G) \setminus \mathbb{R}(\emptyset)$, no set \mathcal{N} of singly-generated continua in $L[G]$ self-collects into $\mathbb{R}(x)$, and thus no nontrivial set of such continua self-constructs.*

Suppose \mathcal{N} did self-collect into $\mathbb{R}(x)$. We will first “zoom in” (if necessary) from all of B to a principal ideal of some ACSA, in order to reduce our situation to the one analyzed in our previous sections.

By Lemma 3.8 let B' be an ACSA of B such that $\mathbb{R}(G \cap B') = \mathbb{R}(x)$. By Lemma 7.4 we may assume that $\|\mathcal{N} \text{ self-collects into } \mathbb{R}(G \cap B')\| \in B'$. We wish to consider B' as our “outermost” algebra, and refer to $G \cap B'$ as G' .

Note that in passing to this (possibly) smaller algebra B' we do not bereave ourselves of all forcing names for \mathcal{N} , since the “ $\mathcal{N} \in L(X)$ ” clause of the definition of self-collection ensures $\mathcal{N} \in L(\mathbb{R}(x)) = L(G \cap B')$, so that there is a B' -name for \mathcal{N} . Thus we assume Θ and χ to be defined relative to B' and to this B' -name.

Finally, in order to have the simplifying situation that “ \mathcal{N} self-collects into $\mathbb{R}(G \cap B')$ ” is forced by 1 (rather than just by some $g \in G$), let us zoom in further to the principal ideal

$$B'' \equiv B' \upharpoonright \|\mathcal{N} \text{ self-collects into } \mathbb{R}(G \cap B')\|,$$

considered as a boolean algebra in its own right.

Now suppose that $D \in \text{ACSAs}(B'')$ and $g \in G'$ witnessed the self-collection of \mathcal{N} into $\mathbb{R}(x)$ in fulfillment of Theorem 8.7. Consider the ACSA D^{+g} and its principal ideal $D^{+g} \upharpoonright g$. By Lemma 9.3, there exists $\phi \in \Phi_0$ that fixes D^{+g} , and is not an identity function. By Lemma 9.5, $\phi \in \Phi$. The existence of such a ϕ contradicts Theorem 8.7. \square

9.2 Cohen forcing does not work in our construction

Theorem 9.7 *No nonempty set of singly-generated continua in a Cohen-forcing extension of L self-collects into a singly-generated continuum.*

The boolean completion of a Cohen-forcing poset satisfies the definition of flexible homogeneity, as we will now show by devoting one lemma to each of that definition’s four clauses; the theorem just stated will then follow from Theorem 9.6.

Definitions. A *Cohen-forcing algebra* is a complete atomless boolean algebra with a countable dense subset. We will see that there is only one such algebra up to isomorphism. Let K denote it.

A *free boolean algebra on κ generators* is the boolean algebra F_κ having a subset $\{g_\alpha : \alpha < \kappa\}$ (called *free generators*) that (1) finitely generates F_κ in the sense that every F_κ -member is the output of some boolean operation on some finite set of g_α ’s, and (2) has “mutually independent” members in the sense that the meet of any finite number of g_α ’s and/or negations thereof will be nonzero (assuming of course that for no α are both g_α and $\neg g_\alpha$ among the conjuncts). Note that F_κ is a complete boolean algebra if and only if κ is finite.

These two defining properties easily entail that whenever A, B are free boolean algebras having sets A_0, B_0 of generators with the same cardinality, any bijection between A_0 and B_0 induces an isomorphism between A and B .

Lemma 9.8 *If K is a Cohen-forcing algebra and $X \subseteq K$ is a (possibly empty) set of free generators for a subalgebra X' that is dense in the complete subalgebra \overline{X} that it completely*

generates (in K), and $\|\overline{X} \geq_G K\| = 0$, then there exists $Y \subseteq K \setminus X$, such that $X \cup Y$ is a set of free generators of a subalgebra of K that is dense therein.

By definition K^+ has a countable dense subset $\{q_n : n \in \omega\}$. We will use this subset to iteratively fix K^+ -members y_n so that the set $Y = \{y_n : n \in \omega\}$ will be as required by the lemma's statement. Let us use K_n to denote the subalgebra of K that will be completely generated by $X \cup \{y_i : i < n\}$. In the notation for one-element expansions from Section 7, we then have

$$\begin{aligned} K_0 &= \overline{X}; \\ K_1 &= \overline{X}^{+y_0}; \\ K_n &= \overline{X}^{+y_0+y_1+\dots+y_{n-1}}. \end{aligned}$$

By repeated invocation of Lemma 7.4 we have $\|K_n =_G \overline{X}\| = 1$, and thus by transitivity (Corollary 3.3), $\|K_n \geq_G K\| = 0$, for all n .

Let our induction hypothesis be that the set $X \cup \{y_i : i < n\}$ is a set of free generators of a subalgebra that is dense in K_n . This clearly holds when $n = 0$.

We now begin the inductive definitions. At stage $n \geq 0$, let m be least such that no finite plus/minus meet from $X \cup \{y_i : i < n\}$ is $\leq q_m$. Some such q_m must exist, lest K_n^+ be dense in K^+ , contradicting $\|K_n \geq_G K\| = 0$. Set $y_n^0 \equiv q_m$ (the superscript "0" here is a secondary index rather than an exponent). If $(y_n^0 \uparrow K_n) = 1$, then $y_n^0 \wedge k > 0$ for all $k \in K_n^+$; for if k were a counterexample, then $1 > \neg k \geq y_n^0$ would contradict $(y_n^0 \uparrow K_n)$'s definition as the *least* K_n -member that is $\geq y_n^0$. So in this case (namely $y_n^0 \uparrow K_n = 1$) we may simply set $y_n \equiv y_n^0$. Otherwise, consider that since $(y_n^0 \uparrow K_n) < 1$, we have $\neg(y_n^0 \uparrow K_n) > 0$. There must exist $k \in K^+$ such that $k \leq \neg(y_n^0 \uparrow K_n)$ but no K_n^+ -member is $\leq k$, lest we have

$$\|K_n \geq_G K\| \geq \neg(y_n^0 \uparrow K_n),$$

violating $\|K_n \geq_G K\| = 0$. So let $y_n^1 \equiv y_n^0 \vee k$ for some such k ; and, in general, iteratively define $y_n^{\alpha+1} \equiv y_n^\alpha \vee k$ for some k satisfying

$$k \leq \neg(y_n^\alpha \uparrow K_n),$$

and no K_n^+ -member is $\leq k$, and define $y_n^\alpha \equiv \bigvee_{\beta < \alpha} y_n^\beta$ for limit ordinals α , until such time as $(y_n^\alpha \uparrow K_n) = 1$. Once this happens, set $y_n \equiv y_n^\alpha$. Note that $y_n \wedge (y_n^0 \uparrow K_n) = q_m$, so that there is a K_{n+1}^+ -member that is $\leq q_m$.

It is then a matter of routine boolean algebra to verify that the induction hypothesis will hold at the next step, and that $Y \equiv \{y_n : n \geq 0\}$ will be as promised. \square

Corollary 9.9 *A Cohen-forcing algebra K has a countable free subalgebra Y' that is dense in K .*

Invoke Lemma 9.8 for K with X empty. \square

Corollary 9.10 *The Cohen-forcing algebra is unique up to isomorphism.*

Say A, B are Cohen-forcing algebras having (by Corollary 9.9) countably infinite free subalgebras A_0, B_0 that are dense in A and B respectively. Any isomorphism between sets of generators for A_0 and B_0 respectively will (as mentioned above) induce an isomorphism between A_0 and B_0 ; and because A_0 and B_0 are dense in A, B respectively, any isomorphism between A_0 and B_0 will extend uniquely to an isomorphism between A and B . \square

Lemma 9.11 *The Cohen-forcing algebra satisfies ACSA-homogeneity (clause 1 of flexible homogeneity).*

This follows from Corollary 9.10 so long as each principal ideal $C \upharpoonright c$ of each ACSA C of a Cohen-forcing algebra K is itself a Cohen-forcing algebra. And this is the case because if X is any countable dense subset of K , then $\{x \upharpoonright C : x \in X \cap (K \upharpoonright c)\}$ is a countable dense subset of $C \upharpoonright c$. \square

Lemma 9.12 *There exists an ACSA C of the Cohen-forcing algebra K satisfying $1 \Vdash C \neq_G K$ (clause 2 of flexible homogeneity).*

Obtain from Corollary 9.9 a countable free subalgebra Y' of K that is dense in K ; let $\{y_n : n \in \omega\}$ be a set of free generators for Y' . Let C_{even} denote the ACSA of K completely generated by $\{y_n : n \text{ even}\}$; and likewise for C_{odd} and $\{y_n : n \text{ odd}\}$. It is then straightforward to show that $1 \Vdash C_{\text{even}} \not\leq_G C_{\text{odd}}$, and therefore $1 \Vdash C_{\text{even}} \neq_G K$. \square

Lemma 9.13 *If $1 \Vdash C <_G K$ then there exists an automorphism of K that is not an identity mapping but whose restriction to C is an identity mapping (clause (3) of flexible homogeneity).*

Use Lemma 9.8 to obtain a set of free generators of a free subalgebra that is dense in C . Call this set X ; and use Lemma 9.8 again, with this X and K , to obtain a set $Y \subseteq K \setminus X$ such that $X \cup Y$ is a set of free generators for a free subalgebra F that is dense in K . The automorphism of $X \cup Y$ obtained by interchanging y and $\neg y$, for some chosen $y \in Y$, and leaving the other generators invariant, induces an automorphism of F , which in turn induces the desired automorphism of K . \square

Lemma 9.14 *Each isomorphism ϕ between any $C, C' \in \text{ACSAs}(K)$ satisfying*

$$1 \Vdash C, C' \neq_G K$$

extends to an automorphism of K (clause (4) of flexible homogeneity).

Fix C, C', ϕ . Obtain X, Y for C as in Lemma 9.13. By isomorphy, X 's image under ϕ will be a set X' of free generators for a free subalgebra dense in C' . Use Lemma 9.8 with this X' and K to obtain Y' . Let θ be any bijection from the countable set Y of generators onto the countable set Y' of generators. It is straightforward to show that $\phi \cup \theta$ induces an automorphism of K . \square

Now Lemmas 9.11, 9.12, 9.13, 9.14 have established that all the clauses of the definition of flexible homogeneity hold for the Cohen-forcing algebra; so Theorem 9.7 is proved.

9.3 Random-real forcing does not work in our construction

A real number that is “random over L ” in the sense established by R. Solovay is one that is interdefinable in a canonical way with a generic filter on the “random-real” algebra, namely the algebra R of Borel subsets of the interval $(0, 1)$ modulo Lebesgue-null sets; see Chapter 15 of [6]. Our goal is to show that R is, like the Cohen algebra, flexibly homogeneous, and so succumbs to Theorem 9.6.

Definitions:

A (strictly positive and normalized, or “probabilistic”) *measure* on a boolean algebra B is a real-valued function μ on B that satisfies

- (i) $\mu(0) = 0$;
- (ii) $\mu(b) > 0$ for all $b \in B^+$;
- (iii) for all pairwise incompatible $b_n, n = 0, 1, \dots$,

$$\mu\left(\bigvee_{n \geq 0} b_n\right) = \sum_{n \geq 0} \mu(b_n);$$

- (iv) $\mu(1) = 1$.

A *measure algebra* is a complete boolean algebra that carries a measure μ . The random-real algebra R is a countably-completely-generated measure algebra when μ is defined to be the Lebesgue measure (recall that R 's members are equivalence classes of sets of reals modulo Lebesgue-null sets).

Two elements $x, y \in R$ are μ -independent if $\mu(x \wedge y) = \mu(x)\mu(y)$. Two subsets $X, Y \subseteq R$ are μ -independent if x, y are μ -independent for every pair $x \in X, y \in Y$.

When $X \subseteq R$, a *plus-minus meet from X* is an element of form

$$\pm x_1 \wedge \pm x_2 \wedge \dots \wedge \pm x_n \tag{8}$$

for some finite $n > 0$, with all $x_i \in X$, where $\pm x$ means either x or $\neg x$, and for no $x \in X$ are both x and $\neg x$ among the conjuncts.

We say $X \subseteq R$ has the “ 2^{-n} property” if, whenever x is a plus-minus meet of n different X -members and/or negations thereof, $\mu(x) = 2^{-n}$.

Lemma 9.15 *There exist subsets $\{e_i : i \in \omega\}, \{o_i : i \in \omega\} \subset R$, each having the 2^{-n} property, whose members are free generators of mutually μ -independent free subalgebras F_{even}, F_{odd} (respectively), such that $\overline{F_{even} \cup F_{odd}} = R$.*

It is well known that R has a subset $\{b_i : i \in \omega\}$ that completely generates R and has the 2^{-n} property. For a concrete example, consider that R is completely generated by the rational subintervals of $(0, 1)$; all these subintervals are completely generated just by those whose endpoints are terminating binary decimals, and these in turn are generated by the subsets of form

$$b_i \equiv \bigcup_{0 \leq n < 2^i} \left(\frac{2n}{2^{i+1}}, \frac{2n+1}{2^{i+1}} \right),$$

where $i \in \omega$. (Note that we have, as is customary in discussions of R , stopped writing “modulo null sets” explicitly.) It is easily checked that $\{b_i : i \in \omega\}$ has the 2^{-n} property.

Given these b_i , we first partition them into two subsets by defining, for all i , $e_i \equiv b_{2i}$ and $o_i \equiv b_{2i+1}$. We claim that these e_i and o_i are as required. Let F_{even}, F_{odd} be the subalgebras of R finitely generated by $\{e_i : i \in \omega\}$ and $\{o_i : i \in \omega\}$. Note that we have ensured $\overline{F_{even} \cup F_{odd}} = R$ because $\{b_i : i \in \omega\}$ completely generates R .

Since $\{e_i\}$ is a subset of $\{b_i\}$, it too has the 2^{-n} property; and from this it easily follows that F_{even} is a free subalgebra for which $\{e_i\}$ is a set of free generators. Similarly for F_{odd} and $\{o_i\}$.

It remains to show that F_{even} and F_{odd} are μ -independent, i.e. that $\mu(e \wedge o) = \mu(e)\mu(o)$ for all $e \in F_{even}, o \in F_{odd}$. Fix e and o . Consider that o is the output of a boolean expression using k -many distinct o_i 's (and no other elements), namely $o_{i_1}, o_{i_2}, \dots, o_{i_k}$. These elements are free generators of a finite boolean subalgebra $O_{fin} \subseteq F_{odd}$ having 2^k atoms, each of form

$$a = \pm o_{i_1} \wedge \pm o_{i_2} \wedge \dots \wedge \pm o_{i_k},$$

(where $\pm x$ means either x or $\neg x$) and each satisfying $\mu(a) = 2^{-k}$. Any element of a finite boolean algebra is uniquely expressible as the join of finitely many atoms; in the case of our $o \in O_{fin}$, say m -many such atoms.

Likewise, e will be a member of some E_{fin} generated by l -many of the e_i 's, and will be the join of finitely many (say n -many) E_{fin} atoms b , each satisfying $\mu(b) = 2^{-l}$. By the 2^{-n} property of the original $\{b_i\}$ ($= \{e_i\} \cup \{o_i\}$), $o \wedge e$ is the join of $m \times n$ many mutually incompatible elements of form $a \wedge b$, each satisfying $\mu(a \wedge b) = 2^{-k-l}$. By additivity of μ on incompatible elements, then,

$$\mu(o \wedge e) = (m \times n)2^{-k-l} = (m \times 2^{-k})(n \times 2^{-l}) = \mu(o) \mu(e). \quad \square$$

Definitions:

Fix subsets $\{o_i : i \in \omega\}, \{e_i : i \in \omega\}, F_{odd}, F_{even}$ as in Lemma 9.15.

Let C_{even} denote $\overline{F_{even}}$ (F_{even} 's completion in R); and $C_{odd} = \overline{F_{odd}}$.

If $\{a_n : n < \omega\}$ is a sequence in R , let $\limsup a_n \equiv \bigwedge_{n=0}^{\infty} \bigvee_{k \geq n} a_k$ and $\liminf a_n \equiv \bigvee_{n=0}^{\infty} \bigwedge_{k \geq n} a_k$. If $\limsup a_n = \liminf a_n = a$, we say that $\{a_n : n < \omega\}$ converges, and let $\lim_n a_n = a$.

We restate the following two claims from [6] more or less verbatim, and without proof.

Fact 9.16 (Exercise 30.1 of [6]:) *If X is a subalgebra of a measure algebra R then the complete subalgebra of R σ -generated by X [which is what \overline{X} denotes] consists of all limits of convergent sequences in X . \square*

Fact 9.17 (Exercise 30.2 of [6]:) *If μ is a measure on R and $a = \lim_n a_n$, then $\mu(a) = \lim_n \mu(a_n)$. \square*

Lemma 9.18 *If X, Y are μ -independent subalgebras of R then their completions $\overline{X}, \overline{Y}$ in R are μ -independent.*

Fix $x \in \overline{X}, y \in \overline{Y}$; we must verify $\mu(x \wedge y) = \mu(x)\mu(y)$. We show that this follows from Facts 9.16 and 9.17. Invoking 9.16, fix $\{x_n\}$ converging to x in the above sense, and $\{y_n\}$ converging to y .

Claim: $\lim_{n \rightarrow \infty} \mu(x \wedge \neg x_n) = 0$. Proof: let \hat{x}_n denote $\bigwedge_{k \geq n} x_k$; then since $x = \liminf x_n$, x is the least upper bound of $\hat{x}_1 \leq \hat{x}_2 \leq \dots$. By μ 's countable additivity we must have $\lim_{n \rightarrow \infty} \mu(x - \hat{x}_n) = 0$. Now if the claim did not hold, there would exist $\epsilon > 0$ such that $\mu(x \wedge \neg x_n) > \epsilon$ for arbitrarily large n ; but then $\mu(x \wedge \neg \hat{x}_n) > \epsilon$ for these same arbitrarily large n , contradiction.

Similarly $\lim_{n \rightarrow \infty} \mu(x_n \wedge \neg x) = 0$, lest $\limsup x_n = x$ fail; and likewise

$$\lim_{n \rightarrow \infty} \mu(y_n \wedge \neg y) = 0; \quad \lim_{n \rightarrow \infty} \mu(y \wedge \neg y_n) = 0.$$

From this it follows by simple boolean manipulation that

$$\lim_{n \rightarrow \infty} \mu(((x_n \wedge y_n) \wedge \neg(x \wedge y)) \vee ((x \wedge y) \wedge \neg(x_n \wedge y_n))) = 0.$$

Thus

$$\lim_{n \rightarrow \infty} |\mu(x_n \wedge y_n) - \mu(x \wedge y)| = 0$$

and

$$\mu(x \wedge y) = \lim_{n \rightarrow \infty} \mu(x_n \wedge y_n) = \lim_{n \rightarrow \infty} (\mu(x_n)\mu(y_n)) = \mu(x)\mu(y),$$

where the last equality holds by Fact 9.17. \square

Corollary 9.19 *C_{even} and C_{odd} are μ -independent. \square*

Lemma 9.20 *R is ACSA-homogeneous (clause (1) of flexible homogeneity).*

Consider any $C \in \text{ACSAs}(R)$. It is easy to verify conditions (i)-(iv) above for μ 's restriction $\mu \upharpoonright C$. The same is true for μ 's restriction to any principal ideal $C \upharpoonright c$ (we satisfy (iv) by defining $\mu'(b) \equiv \mu(b)/\mu(c)$, so that that $C \upharpoonright c$, considered as a boolean algebra in its own right, carries the normalized measure μ'). Thus by Maharam's uniqueness-up-to-isomorphism theorem, $C \upharpoonright c$ is isomorphic to R . \square

Lemma 9.21 $1 \Vdash C_{\text{even}} \neq_G R$ (clause (2) of flexible homogeneity).

Suppose towards a contradiction that $\|C_{\text{even}} =_G R\| > 0$. By Lemma 4.7 there exists $b \in R^+$ such that $b \wedge [R] \subseteq b \wedge [C_{\text{even}}]$, that is,

$$(\forall o \in R)(\exists z \in C_{\text{even}})(b \wedge o = b \wedge z). \quad (9)$$

Recall the free generators o_i of F_{odd} , having the 2^{-n} property, whose completion in R is C_{odd} . Let $n > 0$ be arbitrary, and let M_n denote the set of all plus-minus meets of form

$$a = \pm o_0 \wedge \dots \wedge \pm o_{n-1}$$

satisfying $b \wedge a > 0$. Enumerate M_n as $\{a_1, a_2, \dots, a_k\}$. Note that $i \neq j \Rightarrow a_i \wedge a_j = 0$.

Now for each $a_i \in M_n$, there exists by (9) some $z_i \in C_{\text{even}}$ such that $b \wedge z_i = b \wedge a_i$. For each $a_i \in M_n$ fix some such z_i . If for each i we define

$$z'_i \equiv z_i \wedge \bigwedge_{j \neq i} \neg z_j,$$

we will still have $b \wedge z'_i = b \wedge a_i$. For if $b \wedge z'_i < b \wedge z_i$, then for some $j \neq i$, we have $b \wedge z_i \wedge z_j > 0$. But this contradicts $b \wedge z_i \wedge z_j = b \wedge a_i \wedge a_j \leq a_i \wedge a_j = 0$.

Note that $\{z'_i : i \leq k\}$ is an antichain, and each z'_i belongs to C_{even} . We have:

$$b = (b \wedge a_1) \vee \dots \vee (b \wedge a_k) = (b \wedge a_1 \wedge z'_1) \vee \dots \vee (b \wedge a_k \wedge z'_k) \leq (a_1 \wedge z'_1) \vee \dots \vee (a_k \wedge z'_k).$$

It follows that $\mu(b) \leq \mu((a_1 \wedge z'_1) \vee \dots \vee (a_k \wedge z'_k)) = \mu(a_1 \wedge z'_1) + \dots + \mu(a_k \wedge z'_k)$. Since $a_i \in C_{\text{odd}}$ and $z'_i \in C_{\text{even}}$, Corollary 9.19 implies

$$\mu(b) \leq \mu(a_1)\mu(z'_1) + \dots + \mu(a_k)\mu(z'_k).$$

Now each a_i satisfies $\mu(a_i) = 2^{-n}$ (by the 2^{-n} property); and the z'_i are an antichain so that the mean value of the $\mu(z'_i)$ terms in this expression is $\leq 1/k$. Thus we have

$$\mu(b) \leq k(2^{-n})(1/k) = 2^{-n}.$$

Since n was arbitrary, we have $\mu(b) \leq 2^{-n}$ for all n ; thus $b = 0$, contradicting $b \in R^+$. \square

To show that R satisfies clauses (3) and (4) of flexible homogeneity, we employ tools from the proof of Maharam's theorem in [6], Chapter 30, in particular the next two lemmas.

Lemma 9.22 (see [6], Lemma 30.5) *Let C be a complete subalgebra of R such that $1 \Vdash C \neq_G R$, and let ν be a measure on C such that $\nu(c) \leq \mu(c)$ for all $c \in C$. Then there exists some $d \in R$ such that*

$$\nu(c) = \mu(d \wedge c) \text{ for all } c \in C.$$

We refer to the proof of [6], Lemma 30.5. Note that $1 \Vdash C \neq_G R$ is equivalent to that lemma's assumption on C by our Lemma 4.7. \square

Lemma 9.23 (see [6], Lemma 30.6) *Let C, D be complete subalgebras of R such that $1 \Vdash C, D \neq_G R$; and let f be a μ -preserving isomorphism of C onto D . Then for every $c \in R$ there exist $d \in R$ and a μ -preserving isomorphism $g \supseteq f$ of $\overline{C \cup \{c\}}$, the subalgebra of R generated by C and c , onto $D \cup \{d\}$.*

This is a slight variant of [6], Lemma 30.6. Aside from variable symbol changes, our version is less general in that C and D are required to be subalgebras of the same algebra R , and are considered to carry measures that are restrictions of the same measure μ on the outer algebra; our version is more general in that we require that C, D satisfy $1 \Vdash C, D \neq_G R$ rather than the stronger condition that they be σ -generated by fewer than κ generators, where κ is the “weight” of R (in our case, \aleph_0). [6]’s proof uses the latter condition to justify invoking Lemma 9.22 (or rather 30.5 in [6]); but our weaker condition is manifestly sufficient for this purpose, as it is the condition explicitly required in the statement of Lemma 9.22. \square

Lemma 9.24 *Let $C_0, D_0 \in \text{ACSAs}(R)$ satisfy $1 \Vdash C_0, D_0 \neq_G R$; then each isometric isomorphism $f_0 : C_0 \rightarrow D_0$ extends to an isometric automorphism of R .*

We largely follow the proof of Theorem 30.1 of [6]. Recall our set $\{b_n : n \in \omega\} \subset R$ that completely generates R .

We build $C_0 \subseteq C_1 \subseteq C_2 \subseteq \dots$ and $D_0 \subseteq D_1 \subseteq D_2 \subseteq \dots$, and μ -preserving isomorphisms $f_0 \subseteq f_1 \subseteq f_2 \subseteq \dots$ between the corresponding C_n ’s and D_n ’s, using an inductive back-and-forth construction. At stage $n + 1$ where n is even, we apply Lemma 9.23 to $f_n : C_n \rightarrow D_n$ and $C_{n+1} \equiv \overline{C_n \cup \{b_{n/2}\}}$, obtaining $d_{n+1} \in R$ and the isometric isomorphism $f_{n+1} : C_{n+1} \rightarrow D_{n+1} \equiv \overline{D_n \cup \{d_{n+1}\}}$ such that $f_{n+1} \supseteq f_n$. At stage $n + 1$ for odd n , we apply Lemma 9.23 to $f_n^{-1} : D_n \rightarrow C_n$ and $D_{n+1} \equiv \overline{D_n \cup \{b_{(n-1)/2}\}}$, obtaining $c_{n+1} \in R$ and the isometric isomorphism $f_{n+1}^{-1} : D_{n+1} \rightarrow C_{n+1} \equiv \overline{C_n \cup \{c_{n+1}\}}$ such that $f_{n+1}^{-1} \supseteq f_n^{-1}$.

After iterating n through all of \mathbb{N} we set $C_\infty \equiv \bigcup_n C_n, D_\infty \equiv \bigcup_n D_n, f_\infty \equiv \bigcup_n f_n$, so that f_∞ is an isometric isomorphism $C_\infty \rightarrow D_\infty$. By Fact 9.16, every element of $\overline{C_\infty}$ is the limit of a convergent sequence in C_∞ , and similarly for $\overline{D_\infty}$; so f_∞ extends to a unique isomorphism $f : \overline{C_\infty} \rightarrow \overline{D_\infty}$. Because our construction ensured that $b_n \in C_\infty$ and $b_n \in D_\infty$ for all n , and the b_n jointly generate R , we have $\overline{C_\infty} = \overline{D_\infty} = R$, so f is an automorphism of R that extends f_0 . Fact 9.17 ensures that f is an isometry. \square

Lemma 9.25 *Lemma 9.24 also holds with the word “isometric” stricken (clause (4) of flexible homogeneity).*

Let C_0, D_0, f_0 be as in Lemma 9.24’s statement except that f_0 may not preserve μ . By Maharam’s theorem, there exists an isometric isomorphism $\phi : C_{\text{even}} \rightarrow D_0$. By Lemma 9.24, ϕ extends to an isometric automorphism of R . Let F denote the image of C_{odd} under ϕ . Since $C_{\text{odd}}, C_{\text{even}}$ are μ -independent, D_0 and F must be too, by isometry.

Now define another measure ν on $D_0 \cup F$, by $\nu(f_0(c)) \equiv \mu(c)$ for all $c \in C_0$, and $\nu(a) = \mu(a)$ for all $a \in F$. It follows from the fact that D_0 and F are μ -independent, and jointly generate R , that ν extends to a measure ν' on all of R . Clearly f_0 is an isometric isomorphism with respect to ν' ; thus by Lemma 9.24 it extends to an isometric automorphism of R . \square

Lemma 9.26 *For each C satisfying $1 \Vdash C \neq_G R$ there exists an automorphism on R that is not an identity mapping but whose restriction to C is an identity mapping (clause (3) of flexible homogeneity).*

Invoke Lemma 9.22 with the measure ν on C defined by $\nu(c) \equiv \mu(c)/2$, to obtain $d \in R$ such that $\mu(d \wedge c) = \nu(c) = \mu(c)/2$ for all $c \in C$. Let $C_0 = D_0 = \overline{C \cup \{d\}}$, and let f_0 be the unique μ -preserving isomorphism $C_0 \rightarrow D_0 (= C_0)$ that fixes C and exchanges d and $\neg d$. It is straightforward to verify that this f_0 exists and is unique, using the fact that

$$C_0 = D_0 = \{(c \wedge d) \vee (c' \wedge \neg d) : c, c' \in C\}$$

and the choice of d . Note that C_0 also satisfies $1 \Vdash C_0 \neq_G R$: if some g forced $R \leq_G C_0$ then by Lemma 4.7 at least one of $g \wedge d, g \wedge \neg d$ would force $R \leq_G C$.

Now it suffices to note that f_0 extends to an automorphism on R , by Lemma 9.25. \square

Theorem 9.27 *No nonempty set of singly-generated continua in a random-real forcing extension of L self-collects into a singly-generated continuum.*

This now follows from Theorem 9.6 since we have established that R meets all clauses of the definition of flexible homogeneity (Lemmas 9.20, 9.21, 9.25, and 9.26). \square

10 Prospects For Self-Collection

Theorem 8.7 showed that (under Stipulation 3.11) a forcing algebra B will allow some \mathcal{N} in the generic extension to self-collect into a singly-generated continuum just if B avoids having partial automorphisms of a particular kind. Cohen forcing and random-real forcing algebras do have such partial automorphisms; we showed in Section 9 that this follows

from their satisfaction of the flexible homogeneity property defined there. We now discuss ways that B might avoid this property and so avoid the kind of partial automorphisms that prevent self-collection. In particular we ask how B might violate clause (3) of flexible homogeneity by *rigidly including* subalgebras (a notion introduced in Section 6). The obvious heavy-handed approach would be to use boolean algebras that are themselves rigid, i.e. have no nontrivial automorphisms at all. We will see that this approach can yield sets of continua that self-collect, but that it has a difficulty that may prevent it from providing models of the self-construction axioms. We will then consider the question of whether some B might be nonrigid but still rigidly include subalgebras C ; this approach is subtler, and we currently have no examples of B instantiating it, but we will discuss a potential path to obtaining examples.

A “rigid” approach.

Consider the following construction of P. Hájek, as described by R. Lubarsky in [8]:

Hájek forces over L a sequence of reals $\langle a_n \mid n \in \omega \rangle$ [where a_{n+1} always constructs a_n but not a_{n+2}] which, when coded in some trivial manner into a real α , is a good candidate for the least upper bound of the c [onstructibility]-degrees of the a_n 's. “Good candidate” means that any upper bound which does not collapse \aleph_1 also constructs the sequence. [...]

Hájek's partial order is the ω -step iteration (with finite support) of the Jensen-Johnsbraten forcing [JJ]. The absoluteness comes about because any pair of JJ-generics (over the same ground model satisfying $V = L$) collapses \aleph_1 . Also, a JJ-generic G over L is definable in $L[G]$. So if $\aleph_1^V = \aleph_1^L$ then the sequence of generics is definable as the ω -sequence $\langle a_0, a_1, \dots \rangle$ such that a_{n+1} is the unique set satisfying the appropriate definition over $L[a_n]$.

From this construction we can define a set of distinct continua to serve as our \mathcal{N} :

$$\mathcal{N} = \{\mathbb{R}(a_n) : n \in \omega\}.$$

Now consider $\mathbb{R}(\alpha)$, where α is the real number cited above into which all the a_i have been “coded in some trivial manner” (like interleaving digits). We claim that all five requirements are met for \mathcal{N} to self-collect into $\mathbb{R}(\alpha)$. Clearly $\mathbb{R}(\alpha) \notin \mathcal{N}$. \mathcal{N} is linearly ordered (as $\mathbb{R}(a_0) \subset \mathbb{R}(a_1) \subset \dots$) and so it is directed. By reversing the coding we can reconstruct the sequence of a_i 's from α ; thus $\mathcal{N} \in L(\mathbb{R}(\alpha))$ holds. For requirement (iv), note that any $x \in \mathbb{R}(\alpha)$ that constructs each a_n also constructs the sequence of all the a_n 's (since x does not collapse \aleph_1), and so constructs α ; thus $\mathbb{R}(x)$ cannot be a *proper* subset of $\mathbb{R}(\alpha)$. Finally, the sequence of a_i 's is definable in $L(\bigcup \mathcal{N})$: “... the sequence of generics is definable as the ω -sequence $\langle a_0, a_1, \dots \rangle$ such that a_{n+1} is the unique set satisfying the appropriate definition over $L[a_n]$,” and α is definable wherever the sequence $\langle a_0, a_1, \dots \rangle$ is definable, so indeed $\mathbb{R}(\alpha) \in L(\bigcup \mathcal{N})$.

But this set $\mathcal{N} \cup \{\mathbb{R}(\alpha)\}$ of continua is not self-*constructing*: for any finite $n > 0$, $\mathbb{R}(a_n)$'s predecessors do not self-collect into $\mathbb{R}(a_n)$, since

$$\mathbb{R}\left(\bigcup_{i < n} \mathbb{R}(a_i)\right) = \mathbb{R}(\mathbb{R}(a_{n-1})) = \mathbb{R}(a_{n-1}) \neq \mathbb{R}(a_n).$$

One avenue for seeking a self-constructing set of continua would thus be to modify this construction to obtain a linearly *but also densely* ordered set of continua, from a forcing algebra that, like Hájek's, is free of “bad” partial automorphisms. Our work in this direction has been inconclusive, but we are prepared to say that the task is difficult at best.

The “rigid inclusion” approach.

A more subtle approach would be to seek a B that does have nontrivial partial automorphisms in Φ_0 , but does not cleanly factor into complementary subalgebras in the way that ruled out self-collection in Theorem 9.6, so that $\Phi_1 \neq \Phi_0$. One might try to construct a nonrigid boolean algebra B that *rigidly includes* some ACSA's C , as this was defined in Section 6. We do not know whether such a B exists, but a *-algebraic analog was identified by R. Longo in the 1980s. His paper [7] exhibits inclusions $R \subset M$ of von Neumann algebras such that M 's *-automorphisms are determined by their action on R ; the same then holds of their respective sets of projection operators, considered as complete atomless orthomodular lattices. We hope to investigate whether the same will hold when we pass to the regular open algebras (i.e. boolean completions) of these lattices, considered as complete boolean algebras. We note in closing that forcing with these lattices can be considered a noncommutative version of random-real forcing.

11 Appendix: Review of Relative Constructibility, Basic Facts

The best exposition of constructibility theory is probably the first pair of chapters in K. Devlin's book [3]. The theory is developed in a similar way but more tersely in Chapter 13 of T. Jech's standard set-theory text [6]. We will not reproduce this material here, for, as Devlin writes in his preface, “Constructibility theory is plagued with a large number of extremely detailed and potentially tedious arguments, involving such matters as investigating the exact logical complexity of various notions of set theory.” What we will do here is recall enough of the theory to establish a few basic facts that will be fundamental to our work.

11.1 The Constructible Universe L

The axioms of ZF, Zermelo-Frankel set theory with no Choice axiom, capture faithfully but incompletely the intuitive notion of the hierarchy of all well-founded sets (assuming

there is one unique such hierarchy). In particular ZF gives an incomplete answer to the question of what subsets an infinite set can have, and its answer is still incomplete if a Choice axiom (or its negation) is added. Of course ZF's Power Set axiom ensures that "all" subsets of X themselves form a set, and its Separation axioms ensure that any subset of X that can be defined (by a sentence of the formal language of set theory using finitely many set parameters) will be among "all" these subsets. But ZF leaves open the question of what other subsets, not definable in this way, might also exist.

The parsimonious position that no other subsets should exist is embodied in the *constructible universe* L . L is built by transfinite induction using the function $\text{Def}(X)$, which yields the set of all definable subsets of a given set X . More precisely, $\text{Def}(X)$ is the set of all sets of form

$$\{x \in X : \Psi^X(x, c_1, \dots, c_n)\}$$

where Ψ is a proposition of the language of set theory having one free variable x , and c_1, \dots, c_n are some constants in X , and Ψ^X means Ψ 's relativization to X (any unbounded quantifiers in Ψ are constrained to range over X). L is then defined as the union of all stages L_α of the *constructive hierarchy*:

$$\begin{aligned} L_0 &\equiv \emptyset ; \\ L_{\alpha+1} &\equiv \text{Def}(L_\alpha); \\ L_\alpha &\equiv \bigcup_{\beta < \alpha} L_\beta \text{ when } \alpha \text{ is a limit ordinal}; \\ L &\equiv \bigcup_{\alpha \in \text{Ord}} L_\alpha. \end{aligned}$$

L is a model of ZF ([3], Theorem 1.2; [6], Theorem 13.3) and also of the Axiom of Choice ([3], Theorem 3.8; [6], Theorem 13.18).

The construction of L relativizes quite simply to $L(X)$ for arbitrary sets X : specifically, we "seed" the hierarchy at stage 0 by letting $L_0(X)$ be the transitive closure of $\{X\}$. ($\text{TrCl}(\{X\})$ is taken as the seed, rather than X , so that the resulting model will be transitive, which is a requirement to count as an "inner" model of ZF as that term is generally defined.) We then carry out the construction the same way:

$$\begin{aligned} L_0(X) &\equiv \text{TrCl}(\{X\}); \\ L_{\alpha+1}(X) &\equiv \text{Def}(L_\alpha(X)); \\ L_\alpha(X) &\equiv \bigcup_{\beta < \alpha} L_\beta(X) \text{ for limit } \alpha; \\ L(X) &\equiv \bigcup_{\alpha \in \text{Ord}} L_\alpha(X). \end{aligned}$$

This definition can be found in [6] as Definition 13.24, where it is noted that $L(X)$ is the smallest inner model of ZF having X as a member. To *prove* that L is the smallest inner model ([6], Theorem 13.16) and that $L(X)$ is the smallest inner model having X as a member, we must show that they can be defined in an *absolute* way.

11.2 Absoluteness of L and $L(X)$

Definition. When $\Psi(x, Y)$ is a proposition of the language of set theory having exactly one free variable (x) and exactly one constant symbol (Y), we call Ψ a *Y -basic proposition*; when Y is understood to denote some specific set, we call $\Psi(x, Y)$ a *Y -basic predicate*; and if furthermore the expression

$$\{x : \Psi(x, Y)\}$$

defines the same set in any standard inner ZF model (having Y as a member!) in which it is evaluated, we call $\Psi(x, Y)$ a *Y -absolute basic predicate*.

Fact 11.1 *There is a Y -basic proposition Δ such that when Y is chosen to refer to any set, $\Delta(x, Y)$ is a Y -absolute basic predicate and $\{x : \Delta(x, Y)\}$ equals $\text{Def}(Y)$ as defined above.*

To establish this fact we will simply reference [6], Lemma 13.14 and the initial segment of Chapter 13 leading up to it; or see [3], Lemma 2.4, and the pages leading up to it (where it is the formula $D(v, u)$ that formalizes Def). The terminology used in both references is slightly different from ours, but in both cases the formula's absoluteness is proved by showing that it has a low enough level of logical complexity, in terms of its quantifiers. We will not enter into the details here. \square

The next step in showing the absoluteness of L and $L(Y)$ is to formalize the transfinite construction used in their definitions, and to show that each step preserves absoluteness. We will abstract the proof somewhat so that it can be reused for other hierarchies in our paper.

Lemma 11.2 (Absolute Definability Of Transfinite Hierarchies) *Suppose that Z is some fixed set, that $\Omega(x', Z)$ is a Z -absolute basic predicate for this Z , and that $\Psi(x'', (Y, Z, \beta))$ is a (Y, Z, β) -absolute basic predicate for any ordinal β and any set Y . Then for all ordinals α , the following is a (Z, α) -absolute basic predicate:*

$$\begin{aligned} &(\exists f)(f \text{ is a function;} \\ &\quad \text{dom}(f) = \alpha + 1; \\ &\quad f(0) = \{x' : \Omega(x', Z)\} ; \\ &\quad (\forall \beta < \alpha)(f(\beta + 1) = \{x'' : \Psi(x'', (f(\beta), Z, \beta))\}); \\ &\quad (\forall \beta \leq \alpha)(\beta \text{ is a limit ordinal} \Rightarrow f(\beta) = \bigcup_{\gamma < \beta} f(\gamma)) ; \\ &\quad \text{and } x \in f(\alpha)). \end{aligned}$$

This also holds when $\bigcup_{\gamma < \beta}$ is replaced with $\bigcap_{\gamma < \beta}$.

Note first that a function f defined by all the above clauses except “ $x \in f(\alpha)$ ” will in fact always exist, by ZF’s Axiom Schema of Replacement. Now suppose that for some α , the stated predicate fails to be (Z, α) -absolute. Then some *least* α witnesses this failure. But α cannot be 0 (since $\Omega(x', Z)$ is $(Z, 0)$ -absolute), cannot be a successor ordinal (since then $f(\alpha - 1)$ would be $(Z, \alpha - 1)$ -absolute by the induction hypothesis and $\Psi(x'', (f(\alpha - 1), Z, \alpha - 1))$ is $(f(\alpha - 1), Z, \alpha - 1)$ -absolute), and cannot be a limit ordinal (since the union and intersection operations are absolute). \square

Lemma 11.3 *For any ordinal α and any set Z , L_α is definable by an α -absolute basic predicate, and $L_\alpha(Z)$ is definable by a (Z, α) -absolute basic predicate.*

In both cases we plug propositions $\Omega(\dots)$ and $\Psi(\dots)$ into the template of Lemma 11.2. In the L_α case, let Z be \emptyset , and for $\Omega(x', Z)$ use “ $x' \in \emptyset$ ” or any other always-false assertion, so that L_0 will be the empty set; and for $\Psi(x'', (Y, Z, \beta))$ use the proposition $\Delta(x'', Y)$ that (by Fact 11.1) defines $\text{Def}(Y)$ and is a Y -absolute basic predicate for any Y .

Given any set Z , we obtain the Z -absolute predicate for $L_\alpha(Z)$ the same way, but using the given Z instead of \emptyset , and for $\Omega(x', Z)$ using the assertion “ $x' \in \text{TrCl}(\{Z\})$,” which is clearly definable by a Z -absolute predicate. \square

From this lemma, the absoluteness of L and of $L(Z)$ follows: for any sets x and Z , the assertion $(\exists \alpha)(x \in L_\alpha(Z))$ will be either true in all standard inner ZF models, or false in all such models.

The absoluteness of L allows us to use L -members freely in any Y -absolute basic predicate without affecting its Y -absoluteness. More precisely, if (c_1, \dots, c_n) are an ordered n -tuple of sets in L , and for some Y and Ψ , $\Psi(x, (Y, c_1, \dots, c_n))$ is a (Y, c_1, \dots, c_n) -absolute basic predicate, then

$$\{x : \Psi(x, (Y, c_1, \dots, c_n))\}$$

defines the same set in *any standard inner model having Y as a member* — we need not specify “... having (Y, c_1, \dots, c_n) as a member” since every such model has c_1, \dots, c_n as members. Likewise, if any c_i is a member of $\text{TrCl}(\{Y\})$, we need not specify “... having c_i as a member” once we have specified “having Y as a member.” Therefore we may define an easier-to-use but equivalent absoluteness condition (which is the one given in Section 3.2):

Definition. When y is any set, a y -predicate is a proposition $\Psi(x, c_1, \dots, c_n)$ of the language of set theory, having one free variable x and finitely many constant symbols c_1, \dots, c_n , considered together with fixed referents for the c_i *chosen out of* $L \cup \text{TrCl}(\{y\})$. A y -predicate $\Psi(x, c_1, \dots, c_n)$ is y -absolute if

$$\{x : \Psi(x, c_1, \dots, c_n)\}$$

evaluates to the same set in any standard inner ZF model having y as a member.

Lemma 11.4 *Lemma 11.2 still holds when $\Omega(x', Z)$ is allowed to be a Z -absolute (no longer necessarily basic) predicate $\Omega(x', c_1, \dots, c_n)$, and $\Psi(x'', (Y, Z, \beta))$ is allowed to be a (Y, Z, β) -absolute predicate $\Psi(x'', c'_1, \dots, c'_m)$; the predicate guaranteed by the lemma to be (Z, α) -absolute for all α will now be a (Z, α) -absolute predicate using the c_i 's and c'_j 's as constants. \square*

11.3 More detailed proofs of some constructibility lemmas

Proof of Lemma 3.1: This follows from Lemma 11.3, which states the same thing in terms of absolute *basic* predicates. \square

Proof of Lemma 3.4:

The lemma's claim is: $z \in L(y)$ if and only if $z = \{x : \Psi(x, c_1, \dots, c_n)\}$ for some y -absolute predicate $\Psi(x, c_1, \dots, c_n)$. Note that we consider each c_i 's referent as part of the predicate, and we will freely conflate the c_i 's with their referents.

Clearly the “if” direction holds since $L(y)$ is a standard inner ZF model having y as a member.

For “only if,” suppose $z \in L(y)$. By definition of the $L(y)$ hierarchy, z must appear as member of some lowest level $L_\alpha(y)$, and α cannot be a limit ordinal because limit levels of the hierarchy are just unions of previous levels. If $\alpha = 0$, then by the definition of $L_0(y)$, we have $z \in \text{TrCl}(\{y\})$; thus z itself can legally be used as (the referent of) a constant in a y -absolute predicate. So we can simply let $\Psi(x, z)$ be the proposition “ $x \in z$,” and we have our y -absolute predicate that defines z .

The remaining case is that α is a successor ordinal. In this case, $L_\alpha(y) = \text{Def}(L_{\alpha-1}(y))$, and therefore z must be defined by some formula Ψ relativized to $L_{\alpha-1}(y)$, using only a finite number of $L_{\alpha-1}(y)$ -members c_1, \dots, c_n as constants.

Now we know (Lemma 3.1) that $L_{\alpha-1}(y)$ is itself definable by a y -absolute predicate using constants only for $\alpha - 1$ and $\text{TrCl}(\{y\})$. Say $L_{\alpha-1}(y)$ is so defined by $\Delta(x, \alpha - 1, \text{TrCl}(\{y\}))$. Then the predicate $\Omega(x, \alpha - 1, \text{TrCl}(\{y\}), c_1, \dots, c_n)$ defined by

$$\Omega(x) \iff (\exists M)((\forall v)(v \in M \iff \Delta(v, \alpha - 1, \text{TrCl}(\{y\}))), \text{ and } \Psi^M(x, c_1, \dots, c_n))$$

is a predicate (of free variable x) that defines z and is absolute for standard inner ZF models having $\text{TrCl}(\{y\})$ and all the c_i as members. $\Omega(x)$ need not, however, be a y -predicate (as we have defined the term), since the c_i need not be members of $L \cup \text{TrCl}(\{y\})$.

The key point, though, is that even if (the referent of) some c_i is not a member of $L \cup \text{TrCl}(\{y\})$, it will still be a member of $L(y)$, and will have appeared first at some lower level $L_{\beta+1}(y)$. The task now is to replace any uses of such a constant c_i in Ω with a formula that was used to define its referent at step $\beta + 1$ as a subset of $L_\beta(y)$. We will not formalize this replacement; we will simply note that it can be done, and that the result is a predicate $\Omega'(x)$ that will have a new constant c'_1 denoting β , and finitely many c'_1, \dots, c'_m denoting

members of L_β , as new constant parameters. But the predicate will retain its absoluteness, and will still define z , and will no longer use c_i .

We now wish to iterate this replacement to remove any other c_i 's — and now also any c'_i 's — that are not members of $L \cup \text{TrCl}(\{y\})$. Doing so yields a tree of constants whose referents first appear at stages $L_\beta(y)$, and whose “replacement constants” appear at *lower levels* of the hierarchy. Because constants are replaced by formulas using constants from *lower* ordinal stages, the replacement process must terminate, with a final finite set of constants all in $L \cup \text{TrCl}(\{y\})$, lest there be an infinite descending chain of ordinals. The final replacement predicate that uses all these constants is y -absolute and defines z . \square

Full proof of Lemma 5.1:

The “only if” direction of the lemma is what needs to be proved. Assume $G \in L(\Theta)$ and (invoking Lemma 3.4) that G is definable by a Θ -absolute predicate as

$$G = \{x : \Delta(x, c_1, \dots, c_n)\}.$$

We will now show that there exists a Θ -absolute predicate of the more narrowly-defined form demanded in the lemma's statement, that defines the same set (namely G).

We claim it suffices to show that each c_i used by the given Δ can itself be defined by a Θ -absolute predicate of the more narrowly-defined form. Suppose for all $i \in \{1, \dots, n\}$ that c_i were defined by a Θ -absolute predicate as

$$c_i = \{x : \Delta_i(x, \Theta, G \cap C_i, c'_i)\} \quad (*)$$

for some C_i with $G \cap C_i \in \bigcup \Theta$, and some constant $c'_i \in L$. By clause (ii) of Lemma 4.2, the ACSA C' defined by

$$C' \equiv \overline{C_1 \cup \dots \cup C_n}$$

satisfies $G \cap C' \in \bigcup \Theta$. Thus for all i , since $C_i \in L$, we can alter $\Delta_i(\dots)$ so that it continues to be a Θ -absolute predicate defining c_i , but instead of taking $G \cap C_i$ as a constant parameter, takes $G \cap C'$ and C_i as constant parameters, and uses “ $(G \cap C') \cap C_i$ ” wherever the proposition Δ_i used “ $G \cap C_i$ ”. This yields a Θ -absolute predicate

$$\Delta'_i(x, \Theta, G \cap C', C_i, c'_i)$$

that defines c_i and has the more narrowly-defined form. Once this is done for all the c_i , we can similarly transform the given $\Delta(\dots)$ that defines G , replacing each c_i with the Δ'_i expression that defines it, yielding

$$G = \{x : \Delta'(x, \Theta, G \cap C', C_1, \dots, C_n, c'_1, \dots, c'_n)\},$$

where $\Delta'(\dots)$ has the required narrower form.

To show that each c_i can indeed be defined by some Δ_i as in (*) above, we consider four cases. By definition of Θ -absolute predicate, c_i must be a member of L and/or $\text{TrCl}(\{\Theta\})$.

Case 1 : $c_i \in L$. In this case set $c'_i = c_i$, let C_i be arbitrary, and let $\Delta_i(x, \dots)$ simply be the proposition $x \in c'_i$.

Case 2: $c_i = \Theta$. In this set $c'_i = \emptyset$, let C_i be arbitrary, and let $\Delta_i(x, \dots)$ simply be the proposition $x \in \Theta$.

Case 3: $c_i = \theta(X) \in \Theta$ for some continuum X . By Lemma 4.3, c_i will have a member $G \cap C_i$ for some C_i , and c_i will be the set of all copies of this $G \cap C_i$ on subalgebras in $\text{ACSAs}(B)$, via functions in $\text{ParAut}(B)$. Since $\text{ACSAs}(B)$ and $\text{ParAut}(B)$ are constructible sets, c_i is moreover *absolutely* definable from $G \cap C_i$ this way, as

$$c_i = \{x : \Delta_i(x, \Theta, G \cap C_i, \{\text{ACSAs}(B), \text{ParAut}(B)\})\},$$

for an appropriate proposition Δ_i that formally defines being a copy. Thus we can choose this C_i and Δ_i at our step i , and set $c'_i = \{\text{ACSAs}(B), \text{ParAut}(B)\}$.

Case 4: $c_i = F \in \theta(X) \in \Theta$ for some continuum X and some ultrafilter F . This is handled in essentially the same way as Case 3; here c'_i can be $\{C, \phi\}$, where C is the ACSA on which F is an ultrafilter and ϕ is a particular $\text{ParAut}(B)$ -member that copies $G \cap C_i$ to F .

Note that all the members of filters F that can appear in Case 4 are B -members and hence constructible; thus all other $c_i \in \text{TrCl}(\{\Theta\})$ would have fallen under Case 1, so there are no more cases. \square

12 Appendix: Self-constructing ZF^- models as a solution to Russell's paradox

Since we cannot yet produce a model of the two self-construction axioms, it may seem premature to speculate about whether their conjunction with further axioms is satisfiable. There is, however, a third axiom suggesting a novel foundational approach that would dispatch the Russell-type paradoxes in a robust way. Since interest in this novel approach may serve as extra inducement to find models of the first two axioms, we judge that is worth dedicating a brief appendix to this further axiom.

A member of a self-constructing family \mathcal{F} can be understood as a snapshot taken at a certain point in the organic growth process of the real numbers. With foundational questions in mind, we might consider broadening the scope of this intuitive idea so that it describes a self-constructing mathematical *universe*, each of whose members is the snapshot of *the totality of all mathematical structures* — not just real numbers — that are possible at a given point. In fact, we have already defined “continuum” in terms of just such a totality, namely, the relative constructible hierarchy $L(X)$ that the definition of $\mathbb{R}(X)$ references. Thus the simplest strategy to implement this envisioned broadening would be to keep the

same two axioms for the collection \mathcal{F} of continua, but then turn our attention from \mathcal{F} itself to the collection

$$\mathcal{U} \equiv \{L(X) : X \in \mathcal{F}\}$$

of ZF models constructed from each continuum in \mathcal{F} .

There are two related problems with this strategy; the first one could be called merely technical, but it points directly to the second one, which is fundamental. First, the collection \mathcal{U} just defined cannot be a *set*, because each of its members $L(X)$ contains all the ordinals and so is a proper class. Now, if $L(X)$'s failure to be a set did cause an annoyance in some deduction we would like to make with it, we could presumably use some standard workaround using the fact that “ $y \in L(X)$ ” is first-order expressible using just X as a parameter. But this annoyance would raise the question: are we justified in granting each member $L(X) \in \mathcal{U}$ “all the ordinals” in the first place? No, we claim — this goes against the original intuition we are axiomatizing, namely a collection of mathematical objects that constructs itself without anything given “in advance” or “from outside.” By taking $L(X)$ to embody everything that can be constructed from X we are in fact surreptitiously giving ourselves this structure called “all the ordinals,” for $L(X)$ is defined as the union of levels $L_\alpha(X)$ of a relative constructible hierarchy, where α ranges over all the ordinals. How, why, and whence are all these α 's given?

To address this concern we must grant ourselves only those levels $L_\alpha(X)$ indexed by α 's that are themselves “constructively accessible from X .” There are several approaches we could take to make this precise; here we will stick to just one and leave others unmentioned.

Letting X be a singly generated continuum, as witnessed by $X = \mathbb{R}(x)$ for some real x , and letting $\lambda(X)$ denote the smallest uncountable ordinal in $L(X)$, we consider an ordinal α to be *constructively accessible from X* if $\alpha < \lambda(X)$, and we consider the universe of sets $L^-(X)$ corresponding to the continuum X — the “broadening” of X — to be the union of all the $L_\alpha(x)$ for these ordinals:

$$L^-(X) \equiv L_{\lambda(X)}[x].$$

Note that all sets in $L^-(X)$ are countable and that the continuum X is a subset, but not a member, of $L^-(X)$. Thus $L^-(X)$ violates the powerset axiom and is not a model of ZF. It does, however, satisfy all the other axioms of ZF, which are collectively called ZF^- .

Lemma 12.1 *For any singly-generated continuum X , $L^-(X)$ is well defined (i.e. does not depend on the choice of x that generates X), satisfies ZF^- , and satisfies $\mathbb{R}(X) \subseteq L^-(X)$.*

The key to showing $L_{\lambda(X)}[x] = L_{\lambda(X)}[y]$, for two generators x, y of X , is to prove that y appears at a countable (in $L[x]$) stage α of the $L_\alpha[x]$ hierarchy, and vice versa; and the key to showing this is a generalization, provided in [6] as Lemma 13.24, of the “condensation lemma.” The same lemma is used to show $\mathbb{R}(X) \subseteq L^-(X)$. To prove that $L_{\lambda(X)}[x]$ satisfies each ZF^- axiom, one simply checks that the standard proof that $L[x]$ satisfies it will still work in the absence of the hierarchy's uncountable levels. We omit the details. \square

$L^-(X)$'s failure to satisfy ZF in full does not detract from its suitability to be considered a snapshot of the growth of a self-constructing universe, for the powerset axiom is notoriously non-constructive — that is, it asserts the existence of a set without specifying its members. Nor is the viewpoint unprecedented that all sets are countable in the true mathematical universe. D. Scott, one of the founding parents of the modern (i.e. forcing) era of set theory, writes in his preface to [2]: “Perhaps we would be pushed in the end to say that all sets are *countable* (and that the continuum is not even a set) when at last all cardinals are absolutely destroyed. But really pleasant axioms have not been produced by me or anyone else, and the suggestion remains speculation. A new idea (or point of view) is needed ...” (D. Scott, in the foreword to [2], p. xv). We submit that the present appendix may provide such a point of view.

Since we are now considering a self-collecting set universe rather than just a self-collecting continuum, we must generalize our Foundation axiom, which ensures that real numbers do not slip in unaccountably, to ensure now that no sets of any kind slip in unaccountably. Concretely, this means replacing X, Y, Z , which represent continua in \mathcal{F} in the original version of the axiom, with $L^-(X), L^-(Y), L^-(Z)$, which represent “totalities of mathematical possibilities.” Thus:

Foundation Ax. (general version): $(\forall X \in \mathcal{F})(\forall s \in L^-(X))(\exists Y \in \mathcal{F})$
 $(Y \subseteq X \text{ and } s \in L^-(Y) \text{ and } (\forall Z \in \mathcal{F})(Z \subset Y \Rightarrow s \notin L^-(Z)))$.

We use the term *Bergson history* (recalling the philosophy to which the creation of genuinely new possibilities is central) to denote a family \mathcal{F} of continua that satisfies the Self-Collection Axiom, the general version of the Foundation Axiom, and the following:

Anti-Paradox Ax.: $(\exists X, Y \in \mathcal{F})(L^-(X) \in L^-(Y))$.

Let us emphasize how this last axiom dispatches the set-of-all-sets paradoxes: $L^-(X)$ is the totality of all sets at one point in the growth process that \mathcal{F} embodies, and this totality *becomes* a set, quite harmlessly, at some later point in this process.

We note that the Anti-Paradox Axiom requires that, for some $X \subseteq Y$ in \mathcal{F} , the first uncountable cardinal \aleph_1 of $L(X)$ be collapsed to \aleph_0 in $L(Y)$. In terms of requirements on our forcing construction, this entails that some $C \in \text{ACSAs}(B)$ must be a cardinal-collapsing algebra, and thus that B itself must be such an algebra.

References

- [1] U. Abraham and R. Shore, “The Degrees of Constructibility of Cohen Reals”, Proc. London Math. Soc., Volume s3-53, Issue 2, 1 September 1986, Pages 193–208.

- [2] J. Bell, *Set Theory: Boolean-Valued Models and Independence Proofs*, 3rd ed., Oxford University Press, 2005.
- [3] K. Devlin, *Constructibility*, Springer-Verlag, 1984.
- [4] D. H. Fremlin, *Measure Theory*, in the Cloud.
- [5] P. Hájek, “Some results on degrees of constructibility”, in: Müller G.H., Scott D.S. (eds) *Higher Set Theory. Lecture Notes in Mathematics*, vol 669. Springer, Berlin, Heidelberg, 1978.
- [6] T. Jech, *Set Theory*, 3rd Millennium ed., Springer, 2003.
- [7] R. Longo, “Simple injective subfactors.” *Adv. in Math.*, 63(2):152–171, 1987.
- [8] R. Lubarsky, “There’s No Forcing A Least Upper Bound,” *Proceedings of the American Mathematical Society*, Vol. 105, No. 4 (Apr., 1989), pp. 998-1002.